

New Zealand Glacier Monitoring: End of summer snowline survey 2011

Prepared for New Zealand Ministry of Science and Innovation

May 2012



Authors/Contributors:

A. Willsman
T. Chinn
A. Lorrey

For any information regarding this report please contact:

Andrew Willsman
Technician
Field Team
+64-3-348 8987
Andrew.Willsman@niwa.co.nz

National Institute of Water & Atmospheric Research Ltd
10 Kyle Street
Riccarton
Christchurch 8011
PO Box 8602, Riccarton
Christchurch 8440
New Zealand

Phone +64-3-348 8987
Fax +64-3-348 5548

NIWA Client Report No:	CHC2012-052
Report date:	May 2012
NIWA Project:	CLC)1202-1 Climate Present and Past

© All rights reserved. This publication may not be reproduced or copied in any form without the permission of the copyright owner(s). Such permission is only to be given in accordance with the terms of the client's contract with NIWA. This copyright extends to all forms of copying and any storage of material in any kind of information retrieval system.

Whilst NIWA has used all reasonable endeavours to ensure that the information contained in this document is accurate, NIWA does not give any express or implied warranty as to the completeness of the information contained herein, or that it will be suitable for any purpose(s) other than those specifically contemplated during the Project or agreed by NIWA and the Client.

Contents

Executive summary.....	5
1 Introduction	6
1.1 Glaciers and climate change	7
1.2 The Equilibrium Line Altitude (ELA).....	7
2 Field methods.....	9
2.1 The survey flights.....	9
2.2 March 2011 fieldwork details	9
3 Derivation of the glacier snowline.....	12
3.1 ELAs derived by digitising.....	13
3.2 ELAs derived by interpolation	14
3.3 ELAs derived by snow-patch size when the snowline is obscure	14
3.4 Accuracy of values associated with ELA measurements	14
3.5 Derivation of the Long-term or Steady-state ELAo value.....	14
4 The 2011 snowline results	17
4.1 Photographic coverage of the index glaciers.....	17
4.2 Snowline elevation departures.....	17
4.3 Glacier representitivity	20
4.4 The 2010/2011 glacial climate	20
5 Discussion.....	25
5.1 Ice gain and loss.....	25
6 Acknowledgements	28
7 References.....	28
8 Glossary.....	30
Appendix A Index Glacier ELAs (1977-2011).....	31
Appendix B Index Glacier details	34

Tables


Table 3-1: Derived AARo values for New Zealand glaciers.	15
--	----

Table 4-1: Correlation coefficients of individual snowline departures for each glacier correlated against the mean of all remaining values for the period 1977-201.	21
Table 5-1: Generalised climate for 2010/2011 year and previous 11 years in Southern Alps and inferred glacier snow input.	26
Table 5-2: Vertical gradients of mass balance as measured for Ivory and Tasman Glaciers (mm/m).	27

Figures

Figure 1-1: Location of the 50 index glaciers across the South Island of New Zealand.	6
Figure 1-2: Basic parameters of a glacier.	8
Figure 2-1: Flight paths for the two flights of the 2011 glacier survey.	11
Figure 2-2: Weather maps for Flight one on 12 March (left) and 13 March 2011 (entre), and flight two on 30 March (right).	11
Figure 3-1: Schematic diagrams of stratigraphy at the glacier equilibrium line.	12
Figure 4-1: Histogram plot giving the 2010/2011 snowline departures (raw and response normalised) for each index glacier from the long-term ELAo.	18
Figure 4-2: Mean annual departures (raw and normalised) from the ELAo for all measured glaciers for the entire period of these surveys.	19
Figure 4-3: Pacific-wide mean sea level pressure anomalies (top row) April-June 2010 (left) and July-September 2010 (right), with an expanded view of the pattern over the New Zealand sector (bottom row).	23
Figure 4-4: Pacific-wide mean sea level pressure anomalies (top row) October-December 2010 (left) and January-March 2011 (right) with an expanded view of the pattern over the New Zealand sector (bottom row).	24
Figure 5-1: Raw and Scaled Cumulative Mass Balance Indices for the index Glaciers.	25

Reviewed by



James Renwick

Approved for release by



David Wratt

Executive summary

Glaciers of New Zealand respond to the changing climate, and an integration of these changes over time are recorded by annual aerial surveys. These surveys measure the altitudes of the snowlines of 50 glaciers at the end of summer, as a surrogate for annual glacier mass-balance. The survey has been made in most years since 1977, but rarely do conditions permit all of the 50 index glaciers to be surveyed each year. The surveys are carried out by hand-held oblique photography taken from a light aircraft. Both the absolute and relative positions of the snowlines are recorded and this provides a time series of glacier-climate interaction back to 1977. On average, the latest survey of index glaciers indicated a strong negative mass balance for the 2010/2011 glacier year, i.e. the snowlines, the equilibrium line altitudes (ELAs), were well above their average elevation. There was no winter snow left on 14 of the 50 index glaciers, as the snowline had risen above the maximum elevation of each of these. There was some variation across the Southern Alps this year but this seemed due to variability of index glacier response rather than any geographical trend. Many of the index glaciers also exhibited loss of permanent ice area from the previous glacial year.

The strong negative mass balance (i.e. high ELAs) in the 2010/11 season was the integrated effect of warmer than normal temperatures (with record or near-record highs during a few months at some locations across the country, including the western South Island), above normal sunshine and below normal rainfall across the Southern Alps that largely characterised the ablation season. These conditions were largely induced by prominent high pressure systems that were more frequent than normal. The more frequent 'high' circulation patterns over the South Island that generated the regional climate conditions leading to negative glacier mass balance were associated with one of the strongest La Niña events that has occurred in the last 50 years.

The strong negative mass balance is a change from the near-neutral conditions during the previous year (2009/2010). Prior to this there had been two years of strong negative mass balance (2007/2008, 2008/2009). The last glacial year with significant positive mass balance was 2004/2005, and so the trend over the last 6 glacial years has been neutral to negative. The 2010/2011 negative mass balance is the second largest in the 35 year monitoring period. The only year with a larger negative mass balance departure (1989/1990) has a large degree of uncertainty as it is derived from two indirect measurements. The glaciers have shown a varying pattern of positive (22 years) and negative (13 years) mass balance over the 35 year monitoring period. However, some of the index glaciers with well-defined permanent ice areas have clearly lost ice during the course of the 35 year monitoring period. This mass loss has occurred during large negative mass balance years and has not been replaced during the positive mass balance years.

1 Introduction

The results presented in this report continue an annual glacier/climate photographic monitoring programme, begun in 1977, of the position (altitude) of the end-of-summer snowline on 50 selected index glaciers, arranged in several transects across the Southern Alps (see Figure 1-1).




Figure 1-1: Location of the 50 index glaciers across the South Island of New Zealand.

Between 1977 and 1985, a New Zealand glacier inventory was undertaken from Mount Ruapehu in the North Island at 39° 15' S to southern Fiordland at 45° 57' S. A total of 3144 glaciers were identified for the inventory, where a glacier was defined as a permanent area of snow greater than 0.01 km² (one hectare). In the South Island, average peak summits range from 1850 m in Fiordland to 3000 m in the central Southern Alps and descend to 2000 m in the north-central Southern Alps. To the north east, the Kaikoura ranges reach to over 2700 m, where active rock glaciers have developed under the relatively dry climate. Three North Island volcanic cones reach close to the permanent snowline, but only Mount Ruapehu, with a summit at 2752 m, supports glaciers. Logistics of the glacier overflights do not currently allow the North Island region to be included in annual surveys because of the distances involved.

New Zealand has a humid maritime climate, with the Southern Alps lying across the path of the prevailing westerly winds. Mean annual precipitation rises rapidly from 3000 mm along the narrow western coastal plains to a maximum of 15,000 mm or more in the western part of the Alps close to the Main Divide. From this maximum, precipitation diminishes to about 1000mm in the eastern ranges over less than 30km from the divide. This creates a steep west-east precipitation gradient across the Southern Alps, and the mean altitudes of the glaciers closely follows this gradient (Chinn and Whitehouse 1980).

1.1 Glaciers and climate change

Glacier fluctuations are amongst the clearest signals of climate change, because glaciers are highly sensitive indicators of the earth's surface energy balance. Glaciers register distinct signals of past climate change on scales from decades to millennia. Atmospheric changes are signalled by direct, immediate changes in annual mass balance, which are filtered, and smoothed before they become apparent at the glacier front. Glacier snowline altitudes provide a direct value for annual glacier health and balance. Changes indicated by glacier frontal positions are strongly modulated by glacier response times and dynamics related to climate variability, so determination of annual glacier change using frontal positions is problematic.

1.2 The Equilibrium Line Altitude (ELA)

The winter snowpack normally covers the entire glacier in a wedge shape, with the greatest snow depths near the highest altitudes, tapering to zero at the lower edge (Figure 1-2). This lower margin, or transient snowline, of the snowpack retreats up-glacier as summer melt progresses, until it reaches a maximum altitude for the year at the end of summer (March - April). Located somewhere near the mid-point of the glacier, the end-of-summer snowline indicates where snowfall exactly equals snow loss over the past glacial year (Figure 1-2). This line of demarcation is termed the equilibrium line, which is normally visible as a contrast on the glacier surface between discoloured dust on firn (below the equilibrium line) and the clean snow of the previous winter (above the equilibrium line). It is the altitude of this glacial feature that is defined as the Equilibrium Line Altitude (ELA) by Meier and Post (1962). The ELA is measured each year by the snowline survey. The higher the altitude of the ELA, the less the amount of snow remaining to nourish the glacier after summer melt, indicating a lower mass balance. Conversely, a low annual ELA indicates a large amount of residual snow remaining at the end of summer and a positive mass balance. It follows that there is a unique position for the ELA across the glacier where the volume of snow accumulated over

the past year exactly balances the total volume of snow and ice lost during the year. If the ELA were to lie at (or maintain an average of) this altitude for many years, then there would be no change to the mass or size of the glacier (assuming no other processes are involved, e.g. ice-margin lake development and calving). This unique position of the snowline, which indicates that the glacier is in equilibrium with the climate, is defined as the steady-state Equilibrium Line Altitude (ELAo). A snowline of this altitude will indicate zero change to the balance of the glacier. To avoid confusion between the annual and the long-term ELAs, some authors have also referred to the altitude of the annual glacier snowline as the end-of-summer snowline or EOSS.




Figure 1-2: Basic parameters of a glacier.

A shift in climate can change the glacier mass balance and alter the altitude of the annual ELA. Thus the annual snowline position with respect to the long-term average or steady-state ELAo (see section 3) is used as a surrogate for annual balance changes at each glacier (Chinn, 2005; Chinn, 1995). It is the departure of the glacier snowline from the steady-state ELAo, ie. $ELAo - ELA$ that is reported here. These ELA departure values provide a measure of mass balance changes.

Glacier studies worldwide (e.g. Haeberli et al., 2007) have demonstrated that, on average, the accumulation area is about twice that of the ablation area so that the ELAo lies at an altitude where the ratio of the accumulation area (AAR) to the total glacier area has an average value close to 0.66. For this programme, the steady-state ELAo was initially estimated using AAR values of 0.66 for each glacier, then, as more ELA data were obtained, these approximate values have been progressively adjusted for most index glaciers as outlined in Section 3 below.

2 Field methods

Collection of field data involves flying over the glaciers in a light aircraft to take oblique photographs of the position of the end-of-summer glacier snowlines. A GPS with waypoints has been used since 2007 and this has ensured that the oblique photographs have been taken from a similar position every time. It is worth noting here that the GPS waypoints are not of the glacier snow line itself, but the position in the air above the glacier, to achieve the same oblique photo. The snowlines visible on the photographs have historically been sketched on to a map of each glacier and the resulting accumulation or ablation areas are mapped and measured by digitisation. Since 2008 the area interpretation has been done for selected glaciers by digitally rectifying the oblique photos then mapping the accumulation or ablation areas. The 'snowline altitude' can then be accurately assessed from the glacier area-altitude curve or directly from a digital elevation model.

2.1 The survey flights

On the flights, the "navigator" seated beside the pilot, holds a folder of photographs of each glacier. These photographs are used to closely duplicate the position from where previous photographs were taken. Photographs are taken by small- and medium-format SLR cameras, and since the 2001 flight, in digital format. Data on selected glacier termini, geomorphic features, such as moraines and supra-glacial lakes, are recorded in addition to the index glacier end of summer snowlines. The flights are generally flown between 9,000 ft (2,700 m) and 10,000 ft. (3,000 m). The upper limit is determined by civil aviation regulations.

Significant snow melt continues throughout February and March, but by April there is a high probability that the first winter snowfall in the Southern Alps will have occurred. Experience has shown that although successful surveys have been made in April, there is about a 1 in 4 probability of snow before this time. Every year the challenge of the survey is to measure the highest altitude reached by the rise of the glacier snowline as ablation losses precede, before the first "winter" snowfall. A light fall of fresh snow will conceal the position of the snowline as effectively as a coat of paint. The problem has been standardised by setting the earliest date for the flight at March 1.

'Suitable' weather to fly the entire Southern Alps demands particularly settled conditions. A successful survey cannot be guaranteed as there is also a 1 in 10 probability that there will be no suitable flying weather in the month of March before a fresh snowfall occurs because of the prevailing westerly circulation.

2.2 March 2011 fieldwork details

On Saturday 12th March the first leg of the survey flight departed Wanaka in Flightseeing's Cessna 206 piloted by Andy Woods, with Trevor Chinn, Andrew Willsman (NIWA Dunedin), Tim Kerr (NIWA Christchurch), and Blair Fitzharris. This flight headed south to Mt Larkins and then west to Llawrenny Peaks in Fiordland. Cloud was encountered south of Barrier Peak in Fiordland with Mt Irene and the Merrie Range partially obscured. On the return part of this leg Findlay glacier in the West was completely obscured but the remaining glaciers in the Dart and Matukituki valleys were clear. During Leg 1 some recent fresh snow was apparent on the Fiordland glaciers although it did appear to be variable with clear high snowlines visible on some glaciers and others with snowlines obscured by fresh snow. Leg one ended in Wanaka with a refuel and lunch. After lunch Leg 2 headed north to the Haast

region where further partial cloud was encountered on the Stuart, Lindsay, and Brewster Glaciers. Leg 2 continued up the western side of the Divide with some detours to avoid cloud and photograph glaciers through intermittent holes in the cloud. This leg ended in Greymouth with all but one glacier photographed although some photographs were less than ideal. After an overnight stay in Greymouth, Leg 3 went Northeast via glaciers on Faerie Queen and Mt Ella to the Kaikoura Range. Mt Alarm on the Kaikoura Range had recent fresh snow from a storm the previous week. The return from Mt Alarm was made down the eastern side of the Alps and the index glaciers on this side were clear of cloud with the majority having obvious clear snowlines.

The remainder of March remained fine, warm and settled and the decision was made to repeat the middle section of the survey in late March to capture some more detail on the glaciers. On Wednesday 30th March flight two departed Wanaka in Flightseeing's Cessna 206 piloted by Andy Woods, with Trevor Chinn, and Andrew Lorrey (NIWA Auckland), this flight headed west across the Matukituki, Dart and Arawhata Valleys to the Findlay Glacier. From here the flight turned east and photographed Park Pass, Bryant and Larkins Glaciers. Fine clear conditions were encountered on the remainder of the flight which went north along the western side of the Alps to Mt Whitcombe and the Dainty Glacier, and then returned to Wanaka down the eastern side of the Alps. Compared to flight 1 the snowlines were fractionally higher on most glaciers, and on the eastern glaciers the snowlines were significantly clearer. All index glaciers had snowlines either high on the upper neves, or for some off the top of the glaciers. Figure 2-1 presents an overview of the glacier flight path for 2011. By the end of both flights all the index glaciers had been photographed. The weather situations for the two flights are shown in Figure 2-2.

Subsequent to the second glacier survey flight, the "first winter snowfall" fell in early April and was followed by continuing mild weather which melted most of the new snow. Subsequent mild weather is likely to have removed enough new snow that a successful partial survey could have been made as late as the last week of April.




Figure 2-1: Flight paths for the two flights of the 2011 glacier survey. Flight one completed on 12 and 13 March, and flight two on 30 March.




Figure 2-2: Weather maps for Flight on 12 March (left) and 13 March 2011 (middle), and flight two on 30 March (right). Courtesy of Ross Marsden, MetService New Zealand.

3 Derivation of the glacier snowline

Data in this report are presented as “departures from the ELA₀” which represents “departure of the climate of the year from the mean climate for glacier equilibrium”. Thus an accurate estimate of the position of the ELA₀ is an important part of the programme. Associated glacier parameters used are accumulation area (A_c of Figure 3-1) and ablation area (A_b of Figure 3-1); total glacier area A, or A_b + A_c; and accumulation area ratio (AAR) the ratio of the accumulation area to the entire area of the glacier, A_c/(A_c+A_b).




Figure 3-1: Schematic diagrams of stratigraphy at the glacier equilibrium line. (A) for a year of negative balance (high ELA) and (B) for a year of positive balance (low ELA). Numbers indicate age in years of past firm layers.

At the commencement of this project, the value of the ELA was gained by plotting the observed snowline directly on to a topographic map:-

Map + snowline → ELA

On the average glacier, the long term EOSS (ELA₀) divides the accumulation and ablation areas by a rough ratio of 2:1 (AAR value of 0.6). This position was estimated on small glaciers, and derived from the area-altitude curve on the larger glaciers. As the number of years of data increased, many of the ELA₀ values were subjectively adjusted (see section 3.5 for details of the adjustment methods):-

Est. AAR @ 0.6 — ELA₀, + adjustment → refined ELA₀.

Leading to small annual changes to:-

ELA – ELA₀ → Departure value

The use of a GIS mapping system with digitised areas has added significant accuracy to the data by supplying measured glacier and accumulation or ablation areas:-

Digitised (A_c or A_b) + Area Curve → accurate ELA

Digitised A_c + Glacier Area → accurate AAR

In addition, with a longer time series of annual photographs it is frequently more efficient to directly interpolate ELA values between those of previous years which bracket the current

year's snowline (described below), and more than half of the ELA values have been derived by this method:-

Interpolation	→	direct ELA
Where $ELA - ELA_o$	→	Departure
and $ELA + \text{Area Curve}$	→	AAR

The most significant problems in processing the results are recognising the position of the true end-of-summer snowline (Figure 3-1), especially when;

There has been a recent summer snowfall, which effectively “paints out” the snowline.

The snowline has been only partially recorded due to cloud cover, backlighting or other reasons.

The current snowline is obscure or ambiguous due to limited discolouration of snowpacks of previous years.

A wavy or patchy snowline.

Glacier snowline elevations are normally obtained from detailed mass balance studies where the snowline is mapped as the zero isohyet on the annual mass balance map. Results from glaciers around the world where this technique is used are published in the Glacier Mass Bulletin(WGMS, 2011) In this study the snowlines are derived from oblique aerial photographs by one of “three” methods, depending on the snow cover conditions at the end of summer. Firstly by digitising either the accumulation or ablation area to provide a definitive ELA; secondly by interpolating between photographs of past years where a number of snowlines are close in altitude; and thirdly by comparing the sizes of adjacent snow patches when fresh snow or cloud obscures the snowline of the index glacier. These procedures are given in Chinn (1995) and in Chinn and Salinger (1999).

3.1 ELAs derived by digitising

This method gives the definitive ELA values upon which the interpolation and snow patch methods rely. Here the end-of-summer snowline positions are carefully sketched from the oblique photographs onto detailed base maps of each glacier. The mapped accumulation or ablation zones are then digitised using a GIS programme to accurately measure the areas. From 2009 on, however, some of the oblique digital images are now rectified using the Photogeoref software (Corripio 2004). This software utilised the camera position (GPS input), focal length, image size, NZ Digital Elevation Model, and ground control points (clearly observed features identifiable on the NZMG topographic map sheets) to rectify the image. From the total ablation area for each glacier, the snowline elevation is then read off an area-altitude curve constructed for each glacier. If the accumulation area was measured, the ablation area is found by subtracting the accumulation area from the total glacier area. This method provides a single figure for the glacier ELA for the year, regardless of the shape of the measured area, as it eliminates subjective estimation of the altitude of the snowline. The difference between this altitude and that of the long-term ELA_o indicates the annual mass balance of the glacier. Positive values or high snowline elevation signifies less snow and therefore a negative balance.

3.2 ELAs derived by interpolation

With the many photos now available, for many glaciers it may be more accurate and efficient to obtain the ELA value by interpolation. For each glacier, photos for all years are arranged in increasing area of snow cover (descending order of ELAs). The current year's photograph is then carefully compared and inserted into its appropriate place in the sequence. It has been found that very small differences in snow cover can easily be recognised and that two photos separated by many years can have identical snow coverage. The ELA value is interpolated from the ELA values of the adjacent years. Depending on the similarity of the ELAs, this method frequently places the value of the ELA within a few metres.

3.3 ELAs derived by snow-patch size when the snowline is obscure

Where the true end-of-summer snowlines are obscured by fresh snow, cloud or other reasons, the hidden snowline may be interpolated from the degree of snow cover surrounding the glacier, i.e. the size of the intermittent snow patches. Fresh snow on rock has quite a different appearance from fresh snow on existing snow, and it is commonly possible to "see" the snow-patch outline beneath a light cover of new snow. As in the previous method, photographs of the glacier for all years are arranged in order of increasing snow cover on the glacier, which is also the sequence of the size of the snow-patches surrounding the glacier. The photograph from the latest survey is then slotted into its appropriate place in the snow-patch sequence. The ELA values for this glacier are interpolated from those of adjacent years as described above. This subjective assessment has proved to be surprisingly consistent (see Chinn et al, 2002).

3.4 Accuracy of values associated with ELA measurements

Accuracy of the data is dependent on the accuracy of the digitised A_c and/or A_b areas, glacier area and its area curve. Normally all of these values are measured with a high degree of accuracy provided the glacier maintains a constant size. However many of the smaller glacier have undergone large variations of size when both the area and associated area-altitude should be re-measured each year of change. Associated with small glaciers and area changes are the problems of when the snowline (ELA) rises above the glacier or falls below the glacier terminus. When the ELA falls below the glacier, the ELA can be estimated, but the AAR becomes >1 which has statistical problems. When the ELA rises above the glacier, it is not possible to extrapolate its value and the AAR becomes negative.

3.5 Derivation of the Long-term or Steady-state ELAo value

Glacier studies worldwide have demonstrated that the ELAo lies at an altitude which divides the accumulation area from the ablation area in a ratio of near 2:1, and this ratio has been used extensively for the derivation of paleo-snowline altitudes (Maisch, 1992). The accumulation area ratio (AAR) of accumulation area to the total glacier area has an average value close to 0.6 (Paterson, 1994). However, accumulation ratios can range from about 0.25 to 0.75 (Haeberli, Hoelzle, and Zemp, 2007) with the largest deviations occurring for abnormally shaped glaciers (Table 3-1).

The 2:1 ratio of the accumulation to ablation areas, or AARo, of 0.66 was tested and found to apply to New Zealand glaciers without debris cover, terminal lake or an abnormal shape. The test uses the snowline data from the index glaciers as determined from 29 years of

monitoring given in Hoelzle, et al. (2007). Values of the AARo for each index glacier in Appendix 2 are found from the EOSS vs AAR regression data. They vary considerably with the type of glacier, from 0.09 to 0.84. Closer examination shows that the largest deviations are for glaciers with extensive debris cover, and for those with pro-glacial lakes. The results of Table 3-1 show what happens as these are progressively rejected. The nearer the selection to the morphology of a “normal” glacier, the closer the accumulation area ratio approaches the 0.66 mean. Initial observations of the index glacier AAR values suggest that the most significant of the topographic controls for raising the AAR value (lowering the ELA) appear to be the surface gradients below the ELA and any divergence of ice flow. Conversely any flattening of the glacier tongue lifts the ELA to drive a low AAR value. Surprisingly, parallel flow as is common in ice aprons, does not appear to affect the AAR.

Table 3-1: Derived AARo values for New Zealand glaciers.

Sample	Number	Mean AARo	Std Dev
All index glaciers	50	0.57	0.18
Without (a) rock glaciers	46	0.59	0.16
Without (a) and (b) lakes	39	0.63	0.12
Without, (a),(b), and (c) abnormal shapes	26	0.65	0.10

Mass balance, the specific depths of mass gain or loss over a balance year, do not follow an even change along the glacier profile, and for simplification, mid-latitude glaciers are usually assumed to have a single linear gradient along the longitudinal profile of the glacier. However, for equilibrium, the volume of snow gained during the glacier year equals the volume of the ice lost (using water equivalents). Due to the 2:1 rule used here, the accumulation area is approximately twice the ablation area. Thus for the purposes of this work, it is assumed that the ablation mass balance gradient is twice the accumulation gradient

Values for the long-term ELAo were initially derived by applying an AAR value of 0.6 (Gross, et al. 1976) to the area-altitude curve for each glacier. The ELAo is read off the glacier area curves at 0.4 of the area up from the glacier terminus.

Initially the ELAo values were approximate estimates only, as measured AARs on glaciers in equilibrium vary from 0.5 to 0.75, depending on glacier topography and other factors. The ELAo may then be adjusted using the record of annual ELAs and the annual mean ELA for all index glaciers. The method assumes that the ELAo indicates the snowline position for a zero mass balance, and that ELA changes each year on an individual glacier are linearly proportional to the average change over the entire Southern Alps.

The regression plots for each of the index glaciers are given in Appendix 2, where the annual departures for each glacier are regressed against the annual mean for the Alps without the glacier in question. The correlation and representativeness indicated by these regressions is discussed below. At the zero intercept, which indicates a zero average mass balance for the Alps, the mass balance of the individual glacier should also be zero. The adjustment of the ELAo from its estimated value to a precise value indicated by the dataset is carried out using the constant of the regression equation.

The slope of the regression line indicates the character of the response of the glacier to the climate, within the constraints of the assumptions of linearity. The average climate response is given by the mean for the Southern Alps, so that any deviation from a 45° slope is an indicator of the individual characteristics of the glacier. Since each year's climate is thought to be similar over all of the Southern Alps, with the noted exception of the Kaikoura Ranges, which lies in a distinctly different climate district (Mullan 1998, Mullan and Thompson 2006)), the regression slope changes must represent a topographic signal in the ELA values. Similarly, the range of the highest and lowest ELA values is also influenced by the glacier topography. The significance of these properties has yet to be analysed and warrants further investigation.

4 The 2011 snowline results

4.1 Photographic coverage of the index glaciers

During flight one on the 12th March some of the glaciers in the west and south were either partially, or in the case of Findlay Glacier completely, obscured by cloud. Glaciers to the east of the Main Divide on flight one remained cloud-free. The second flight on 30th March was completed in fine cloud-free conditions so many of the earlier cloud-covered index glaciers in the central section of the Southern Alps were photographed on this flight. A low resolution image of each index glacier photograph is in Appendix 2.

The photographs on flight 1 were taken with a Nikon D200 digital camera (sensor size = 23.6 x 15.8 mm, effective pixels = 10 MPix) linked to a Garmin GPSmap 60C_{sx} with an external aerial on the windscreen of the plane. Each digital image has the GPS location (latitude and longitude WGS84 datum) and focal length embedded in the EXIF information part of the digital photograph. The GPS used on the survey was set to the NZMG projection system and was displaying these values on screen, but was outputting WGS84 latitudes and longitudes through the serial interface to the digital camera. These photographs are available as compressed jpegs, or raw Nikon NEF file formats.

The digital photographs on flight 2 were taken with a Pentax K100D (sensor size = 23.5 x 15.7 mm), this camera did not have a GPS connected to it.

The snowlines were at high elevations on nearly all index glaciers, often there was only remnants of the winter snow left on the upper neves. Fourteen of the fifty index glaciers had no winter snow left on them as the snowline elevation was above the maximum height of the glacier, on these only the previous season's firn and ice remained. Three (Jack, Chancellor, and Jalf) of these fourteen glaciers had snowlines at higher elevations on adjacent glaciers, and these adjacent snowlines were used to extrapolate the respective snowline elevations. During flight one a light dusting of fresh snow from the previous week was observed on the southern glaciers, and these glaciers were not re-photographed on flight two so for these the snow patch method was utilised to determine the snowlines.

4.2 Snowline elevation departures

Monitoring results for the 50 index glaciers for the 2010/2011 glacial year, together with the means for all measured years from 1977 to 2011 are shown in Figure 4-1 and 4-2 respectively.

The individual departure response of each index glacier varies due to topographic factors (discussed in section 3.5) and this variation can be normalised to consistent values by applying the slope from the regression between the annual departure and the annual mean alps value (regression slope values in Appendix 2). The derived slope values are considered to be reasonably valid as there are now a considerable number of observations, with 40 index glaciers having 26 or more annual observations and the least number of annual observations is 17. Figure 4-1 presents the raw and normalised departure values from ELA_o for the index glaciers over the 2010/2011 glacial year. The effect of normalising is minimal for most of the index glaciers which have a near 1:1 ratio response to the Alps mean. Normalising does have the effect of reducing the magnitude of the offset for some of the glaciers with large offset or range of ELA values. It is assumed that the most inconsistent

ranges in ELA values were those for “snow-patch” glaciers which do not have the conventional glacier elevation range, nor an ELA position. Up until now these values have been assigned as theoretical values and application of normalising has brought these difficult sites into line with normal glaciers. For example the 2011 departures for Mt Larkins and the Douglas Glacier are almost halved after the topographic normalising is applied. This normalising for each glacier is especially important when aggregating the results for the full length of record for the whole of Southern Alps, as it accounts for the irregular sample size and the impact of individual glaciers with high sensitivities.




Figure 4-1: Histogram plot giving the 2010/2011 snowline departures (raw and response normalised) for each index glacier from the long-term ELAo.

All of the individual index glaciers for 2010/2011 have ELA departures that are well above the long-term mean ELAo position (Figure 4-1). Some variability exists between index glaciers after normalisation of departures and this is due to a possible combination of geographical variability and measurement uncertainty. In Figure 4-1 there are 11 index glacier observations missing as the snowline was above the maximum elevation of the respective glaciers. Three glaciers with snowlines above the maximum glacial extent have elevations determined from snowlines on adjacent higher permanent ice fields (Jalf, Jack, and Chancellor).

The raw snowline departure for this year (2010/2011) averaged 208 m above the long-term mean ELAo position for the 39 glaciers that had observations. The average of the normalised

2010/2011 observations was the same 208m above the long-term mean ELAo position (Figure 4-2).




Figure 4-2: Mean annual departures (raw and normalised) from the ELAo for all measured glaciers for the entire period of these surveys.

Annual data for all measured glacier ELA departures from the long-term ELAo from 1977 to 2011 are given in the matrix of Appendix 1 and are presented in Figure 4-2. The 2010/2011 departure of 208m above the long term mean ELAo position is a change from the near neutral response of the 2009/2010 glacial year. The 2010/11 departure is the second highest departure recorded in the 35 years of record, only 1989/1990 had a larger departure and this value has a large degree of uncertainty as it is based on two indirect observations correlated to the index glaciers. The large departure during 2010/2011 indicates high snowlines with very small amounts of the winter snowpack remaining on the index glaciers and hence a large loss of snow and ice from the glaciers. The last season of large departures below the ELAo position indicating positive mass balance was the 2004/2005 year, since then annual departures have tended to above or around the ELAo position indicating loss over the last 6 years. This trend of loss of snow and ice is apparent at many of the index glaciers by loss of permanent ice area and fragmentation of many of the small glaciers.

The mean annual departures are presented in Figure 4-2 as raw and normalised amounts, and interestingly the results are similar as the effect of the outliers is very small. The largest change due to normalisation occurred during 1991/1992 where the mean annual departure is reduced by 22m from -125m to -103m, due to a small sample size of 15 index glaciers that contained some strongly sensitive glaciers.

Full data records for each individual index glacier are given in Appendix 2. Missing values are years of no data for the particular glacier.

Appendix 2 gives a data table, map and histograms of all measured snowline fluctuation histories as metres of departure from the steady-state ELAo for each index glacier. On the annual departure plots, missing values are years of no survey. The data table provides essential glacier data, snowline data statistics and a table of all measurements and immediate derived values. Photographs of each glacier are available in the 1999 Report, (Chinn and Salinger, 1999) and are reproduced in low resolution in this report.

4.3 Glacier representitivity

The “representativeness” of each glacier as an indicator of the overall annual climate of the Southern Alps is indicated by how well the annual values for an individual glacier correlates with the mean value of the 50 index glaciers over the Southern Alps. Correlation coefficients of individual snowline departures for each glacier correlated against the mean of all remaining values for each year are given in Table 4-1. The correlation plots for each glacier are given in the Appendices. The correlations give a surprising result where representativeness appears to be independent of size, gradient or topography. The high correlations indicate that the ELA surface of individual glaciers have a strong relationship with the “mean ELA” over the whole alpine range. This follows the finding of Clare et al. (2004) where it was demonstrated that the entire Southern Alps behaves as a single climatic unit. However the consistently low correlation of the Kaikoura Range glacier suggests that the behaviour of this range is that of a separate climate zone, while it is assumed that accumulation on the low correlation Langdale glacier is dominated by wind redistribution.

4.4 The 2010/2011 glacial climate

The climate diagnostics for the 2010/2011 glacial year saw conditions less conducive to accumulation (reduced precipitation) during the whole of the austral cool season (winter/spring). Circulation patterns that supported increased ablation (warmer-than-normal temperatures) during the austral warm season (summer/autumn) also existed. More frequent anticyclones (‘highs’) were located over and to the east of the North Island during April, resulting in a very dry month for upper South Island, but enhanced northwest winds affected the lower South Island, producing extraordinarily high rainfall in the southwestern regions including Fiordland, south Westland and parts of Southland, with more than double normal rainfall experienced there. Temperatures were mostly above normal for that month. May was a stormy month overall, with more lows than normal located in the Tasman Sea, and higher pressures east of the Chatham Islands. This pattern produced more northeasterly winds than normal over New Zealand, leading to very sunny and dry conditions on the West Coast, Fiordland and Southland, with less than 50 % of normal rainfall recorded. June 2010 was a very wet month for most regions of the country, with active lows originating over the Tasman Sea affecting the country during both the first and last week of the month, however, below

Table 4-1: Correlation coefficients of individual snowline departures for each glacier correlated against the mean of all remaining values for the period 1977-201.

GLACIER	Correlation Coefficient
BARRIER PK.	0.93
JALF GL	0.92
FINDLAY GL.	0.91
VERTEBRAE #25	0.90
MT. STUART	0.90
LLAWRENNY PKS.	0.89
SIEGE GL	0.89
THURNEYSON GL	0.89
BRYANT GL.	0.89
JACKSON GL.	0.88
MT. BUTLER	0.88
CAROLINE PK.	0.88
MT. ST. MARY	0.88
LINDSAY GL	0.87
MARMADUKE GL.	0.87
CHANCELLOR DOME	0.87
SALISBURY GL	0.87
TASMAN GL.	0.87
MT. CARRINGTON	0.87
MT. LARKINS	0.86
KEA GL	0.86
MT McKENZIE	0.86
MT. GENDARME	0.86
MT. FRANKLIN	0.85
JASPUR GL	0.84
FOG PK	0.84
DAINTY GL	0.84
BREWSTER GL.	0.84
MERRIE RA.	0.83
JACK GL.	0.83
MT. IRENE	0.83
VERTEBRAE #12	0.83
MT. ELLA	0.82
AILSA MTS.	0.82
DOUGLAS GL	0.81
PARK PASS GL.	0.80
MT. AVOCA	0.80
MT. GUNN	0.80
BLAIR GL.	0.80
BROWNING RA	0.79
MT. WILSON	0.77
MT. CARIA	0.76
ROLLESTON GL.	0.76
GLENMARY GL.	0.75
RIDGE GL.	0.74
RETREAT GL	0.72
MT FAERIE QUEENE	0.71
LANGDALE GL.	0.70
SNOWY CK	0.66
KAIKOURA RA	0.59

normal rainfall (50 - 80 % of normal rainfall) was recorded on the West Coast, in Fiordland, and across much of Southland. Temperatures were near average across the Southern Alps for May and June, largely speaking, and this three month period was largely drier than normal.

The following three months had contrasting precipitation and temperature patterns. July 2010 was characterised by higher than normal pressures over the country that brought clear skies, dry conditions, warmer afternoons but colder mornings to many regions. Southeast winds frequently affected New Zealand over the month, and the effects of these were very clear – western regions of the country were extremely sunny and very dry. For the following month, lower pressures than normal over the Tasman Sea and New Zealand brought frequent northerly air streams to the North Island during August, with more easterly winds over the South Island. It was an extremely wet August across most of the country. Mean temperatures were above average (between 0.5°C and 1.2°C above average) across almost all regions of New Zealand. It was very cloudy over much of the country. September 2010 was characterised by extremely low pressures over New Zealand, bringing wild westerly winds. The effect of the stronger-than-normal westerly winds during September was very clear – rainfall was record high or well above average, and sunshine hours were well below average, in western areas of both islands. It was also much cooler than usual in the west and south of the South Island. Extremely warm temperatures affected the country at both the start and end of the month – but an intense southwesterly event from the 17th until 24th brought snow to very low levels in the far south, and record low temperatures there.

New Zealand temperatures during the Austral warm season (summer and autumn) were in stark contrast to those in the preceding three months. Many stations registered above normal temperatures in addition to below normal precipitation, which contributed to mass balance deficit over much of the Southern Alps for the 2010-11 season. Anticyclones ('highs') dominated New Zealand's climate during October and November 2010, following a rather stormy September. Mean sea level pressures for the season overall were at least 4hPa higher than usual over the Tasman Sea and New Zealand. The prevalence of highs during spring produced a sunny season in many regions, and a much drier than normal spring for South Canterbury, the Lakes District, central Otago, and parts of Fiordland. By the end of spring, the dry conditions, elevated sunshine and above average temperatures had resulted in severe soil moisture deficits (more than 130 mm of deficit) in parts of the Lakes District and central Otago, with significant soil moisture deficits (more than 110 mm of deficit) in many other parts of Canterbury and Otago. Spring seasonal rainfall was below normal (between 50 - 80 % of normal) in south Canterbury, the Lakes District, and parts of Fiordland and central Otago during late 2010. The New Zealand national average temperature for spring was 12.4°C (0.3°C above the 1971–2000 spring average). The regional and national-scale conditions were largely induced by circulation anomalies linked to a building La Niña in the equatorial Pacific. In addition, The Southern Annular Mode (SAM), a climate pattern affecting the westerly wind strength and location over and to the south of the country, was strongly positive overall at the end of 2010. The positive SAM contributed to the poleward migration of the westerly jet, and collaborated with La Niña to produce more frequent anticyclones over New Zealand.

On the transition to summer, La Niña was fully developed while coupling between the atmosphere and ocean was strong. This element of ENSO development is particularly

important for dictating the persistence of characteristic weather patterns associated with La Niña during summer, which typically consist in the New Zealand sector of either more frequent anticyclones as a whole across the country, or more frequent northerly and easterly quarter windflow with blocking highs situated east of the country (Kidson, 2000). The types of 'blocking' patterns all usually serve to reduce precipitation in the major glacier catchments on the South Island, and promote above normal temperatures during summer.




Figure 4-3: Pacific-wide mean sea level pressure anomalies (top row) April-June 2010 (left) and July-September 2010 (right), with an expanded view of the pattern over the New Zealand sector (bottom row).




Figure 4-4: Pacific-wide mean sea level pressure anomalies (top row) October-December 2010 (left) and January-March 2011 (right) with an expanded view of the pattern over the New Zealand sector (bottom row).

5 Discussion

5.1 Ice gain and loss

Glaciers accumulate the mass changes of net annual balance variations over years to decades. These effects of yearly climate variations are delayed and distorted before being delivered to the terminus after individual glacier response times have elapsed. The index glaciers record the annual climate-related mass gains and losses with some degree of accuracy as the majority of these glaciers are small and steep with relatively fast response times, and have areas that are in equilibrium with the climate of recent decades. The large valley glaciers with long, near 100 yr response times have large surface areas inherited in a previous climate, and all are in a state of ongoing recession mainly by downwasting of their debris-covered trunks, but recently ice loss has accelerated by the formation of pro-glacial lakes. For these glaciers, the ELA changes measure ONLY the mass balances of a smaller area that would be in equilibrium with the present climate. Ice mass changes in these glaciers are accounted for in Chinn et al. (2011, submitted).

To assess the mass changes in response to climate fluctuations a cumulative plot of the “mass balance indices” (MBI) is presented in Figure 5-1. The raw MBI is the inverse of the mean annual departure value and this represents the mean departure from steady state (ELA₀). Snowline departure changes with negative ELA (i.e. lower snowline) result in positive MBI. The reliability of use of the ELA as an indicator of mass balance change has been investigated by Chinn et al (2005), where the *r* (correlation) values between the ELA and measured mass balance had an average of 0.9 with a standard deviation of 0.07.




Figure 5-1: Raw and Scaled Cumulative Mass Balance Indices for the index Glaciers.

The annual trends in the raw MBI as shown in Figure 5-1 agree with the generalised climates given in Table 5-1. However, the cumulative total of the raw MBI on the plot does not agree with the obvious ice volume decrease at most of the index glaciers. Over the course of the 35 year monitoring period, there has been permanent ice loss during large negative mass balance years that has not been recovered after a cycle of positive mass balance years. A negative mass balance year has a greater impact on ice volume than a positive mass balance year for MBIs of the same magnitude. This is due to ablation rates typically being almost twice the accumulation rates on most glaciers.

Table 5-1: Generalised climate for 2010/2011 year and previous 11 years in Southern Alps and inferred glacier snow input.

Glacier year	Generalised Climate	Inferred glacier snow input
1997/1998	Higher frequency of anticyclones and westerly winds over the south, southerlies further north. Temperatures 0.2 C below normal, but a very warm summer.	Average
1998/1999	Stronger westerly and northwesterly winds over New Zealand, temperatures 0.8°C above average, with above normal precipitation on the West Coast.	Low
1999/2000	Very anticyclonic, with weaker westerlies than normal. Temperatures 0.7 C above normal, and rainfall slightly below normal.	Low
2000/2001	More northwesterlies over the South Island, temperatures 0.2 C above normal. Rainfall close to average.	Average-High
2001/2002	Higher than normal pressures and more easterlies over the South Island, temperatures 0.3°C above normal, well below average rainfall.	Low
2002/2003	Persistent westerlies and southwesterlies over New Zealand. Cooler spring, rainfall slightly above average in the west and south.	Average-High
2003/2004	More cyclonic westerlies and south westerlies over the South Island from September.	Higher
2004/2005	Cool westerlies during autumn and early winter (temperatures 0.4°C below normal), then strong cold cyclonic southwesterlies through to December (temperatures 0.6°C below normal), precipitation overall close to average.	High
2005/2006	More anticyclones and mild westerlies and northwesterlies during autumn and winter (winter temperatures 0.7°C above normal), then more frequent southeasterlies during spring bringing low precipitation.	Low
2006/2007	More southwesterlies June – November bringing increased accumulation, but then anticyclones and south easterlies December – March with low precipitation and increased ablation. Temperatures near average overall, and precipitation above average.	Average
2007/2008	Variable circulation April – August with little accumulation. From September on, mainly easterly circulation, and especially warm (1.0°C above normal) from December with low precipitation and much increased ablation.	Very low
2008/2009	Northerly and easterly quadrant flow anomalies related to La Niña, with associated normal to above normal temperatures, except during Spring. Below normal precipitation during late Winter and Summer.	Low
2009/2010	Highly variable year with regard to temperature and precipitation swings within and between seasons, particularly for winter 2009 and summer 2010. More frequent southwesterly flow as a result of El Nino development from spring was opposed to record high temperatures in August 2009 and February 2010.	Average
2010/2011	One of the strongest La Niñas in the last 50 years induced more frequent anticyclones (settled conditions) than normal across New Zealand during	Very Low

summer and autumn. 'Highs' were prominent across the South Island during the ablation season, with warmer than normal temperatures as a whole, elevated sunshine hours, and reduced rainfall.

To account for this difference a scaled mass balance index is also shown in Figure 5-1 where negative mass balance years have been scaled by 1.92 and a positive mass balance year departures 1 (i.e., left unchanged). The scale factors are averages of published mass balance gradient rates (Table 5-2) from studies on the Tasman (Anderton, 1975) and Ivory Glaciers (Anderton and Chinn, 1978). (for the period 1966-1975). The assumption is that these averaged mass balance gradients apply to the averaged annual departure value for the index glaciers. To examine the sensitivity of this assumption, and its impact on the cumulative scaled mass balance we have used a range of values for the negative mass balance of 1 (equal to the raw MBI), 1.5, 2 and 3 as shown in Figure 5-2.

Table 5-2: Vertical gradients of mass balance as measured for Ivory and Tasman Glaciers (mm/m). * is an estimated value.

Zone year	Ivory Glacier Ablation	Ivory Glacier Accumulation	Tasman Glacier Ablation	Tasman Glacier Accumulation	Both Glaciers Ablation	Both Glaciers Accumulation
1966-67				6		
1967-68				8.6		
1968-69				7.5		
1969-70	26.3			5.4		
1970-71	35.5		11.7	5		
1971-72	20.8		13.7	7.4		
1972-73	28.9		12.4	10.1		
1973-74	28		14.3	7		
1974-75	15.4		10.4	6.6		
Mean	25.8	12.9*	12.5	7.1	19.2	10

The trend in permanent ice area in many of the index glaciers agrees with the cumulative scaled (1.92) mass balance index rather than the cumulative raw mass balance index which does not account for the mass balance gradient. Further work needs to be done to determine what scaling is appropriate for the index glaciers.

6 Acknowledgements

This research was carried out under Contract C01X0701 with the Foundation for Research, Science and Technology. Tim Kerr kindly provided Andrew with ARC GIS training and passed on his knowledge in the use of the Photogeoref software for rectifying oblique aerial photographs. James Renwick is thanked for providing reanalysis data on three-monthly circulation patterns. Large parts of the climate text are drawn from the monthly and seasonal climate summaries produced by the National Climate Centre (Georgina Griffiths) who are thanked for that resource.

7 References

- Anderton, P.W. (1975). Tasman Glacier 1971-73. Hydrological Research. Annual Report No. 33. Ministry of Works and Development for the National Water and Soil Conservation Organisation, Wellington, New Zealand. 28p
- Anderton, P.W.; Chinn, T.J.H. (1978). Ivory Glacier, New Zealand, an IHD basin study. *Journal of Glaciology* 20(82): 67–84.
- Chinn, T.J.H. (1995). Glacier fluctuations in the Southern Alps of New Zealand determined from snowline elevations. *Arctic and Alpine Research* 27(2): 187—197.
- Chinn, T.J.H.; Heydenrych, C.; Salinger, M.J. (2002). New Zealand Glacier Snowline Survey 2002. NIWA Client Report AKL2003- 023. 20 p. National Institute of Water and Atmosphere, Auckland.
- Chinn, T.J.H.; Heydenrych, C.; Salinger, M.J. (2005). Use of the ELA as a practical method of monitoring glacier response to climate in the New Zealand's Southern Alps. *Journal of Glaciology* 51(172): 85-95.
- Chinn, T.J.H.; Salinger, M.J. (1999). New Zealand Glacier Snowline Survey 1999. NIWA Technical Report 68. 119 p. National Institute of Water and Atmosphere, Wellington.
- Chinn, T.J.H.; Salinger, M.J.; Fitzharris, B.B.; Willsman, A.P. (2009). Annual ice volume changes 1976-2005 for the New Zealand Southern Alps, (submitted).
- Chinn, T.J.H.; Whitehouse, I.E. (1980). Glacier snow line variations in the Southern Alps, New Zealand. pp. 219—228. *In: World Glacier Inventory. International Association of Hydrological Sciences Publication No. 126.*
- Clare, G.R.; Fitzharris, B.B.; Chinn, T.J.H.; Salinger, M.J. (2002). Interannual variations in end-of-summer-snowlines of the Southern Alps of New Zealand, and relationships with Southern Hemisphere atmospheric circulation and sea surface temperature patterns. *International Journal. of Climatology* 22: 107-120.
- Corripio J.G. (2004). Snow surface albedo estimation using terrestrial photography. *International Journal of Remote Sensing Vol 25, Issue 24: 5705-5729.*

- Gross, G.; Kerscher, H.; Patzelt, G. (1976). Methodische Untersuchungen über die Schneegrenze in alpinen Gletschergebieten. *Zeitschrift für Gletscherkunde und Glazialgeologie* 12: 223-251.
- Haeberli, W.; Hoelzle, M.; Zemp, M. (2007). Glacier Mass Balance Bulletin, Bulletin No. 9 (2004-2005) WGMS, 99 p. Staffel Druck Press, Zurich..
- Hoelzle, M.; Chinn, T.J.H.; Stumm, D.; Paul, F.; Zemp, M.; Haeberli, W. (2007). The application of inventory data for estimating characteristics of and regional past climate-change effects on mountain glaciers: a comparison between the European Alps and the New Zealand Alps. *Global and Planetary Change* 56: 69-82.
- Kidson, J.W. (2000). An analysis of New Zealand synoptic types and their use in defining weather regimes. *International Journal of Climatology* 20: 299-316.
- Maisch, M. (1992). Die Gletscher Graubuündens. Geographisches Institut der Universität Zürich.
- Meier, M.F.; Post, A.S. (1962). Recent variations in mass net budgets of glaciers in western North America. pp. 63—77. *In: Proceedings of Obergurgl Symposium, International Association of Hydrological Sciences Publication 58.*
- Mullan, A. B., 1998: Southern Hemisphere sea surface temperatures and their contemporary and lag association with New Zealand temperature and precipitation. *Int. J. Climatol.*, 18, 817-840.
- Mullan, A. B., and C. S. Thompson, 2006: Analogue forecasting of New Zealand climate anomalies. *Int. J. Climatol.*, 26, 485-504.
- Paterson, W.S.B. (1994). *The Physics of Glaciers*. Third edition, Oxford, Pergamon Press. 480 p.
- WGMS 2011. Glacier Mass Balance Bulletin No.11 (2008-2009). Zemp, M., Nussbaumer, S.U., Gartner-Roer, I., Hoelzle, M., Paul, F., and Haeberli, W. (eds.), World Glacier Monitoring Service, Zurich, Switzerland, 102pp.

8 Glossary

Ablation All processes by which snow and ice are lost from a glacier

Accumulation All processes by which snow and ice are added to a glacier.

Accumulation Area Ratio (AAR) The ratio of the accumulation area above the equilibrium line, to the entire area of the glacier.

Departure of the ELA The elevation difference between the long-term ELAo and the annual snowline altitude. Positive departures mean a higher snowline and therefore a negative mass balance.

ELA The mean altitude of the snowline or equilibrium line across a glacier at the end of summer.

ELAo The long-term or steady-state altitude of the ELA which will maintain the glacier in equilibrium with the climate.

Mass balance index The negative of the ELA departure value. This gives values for annual changes with the same sign as the mass balance changes.

Shaded cells Areas which have been measured by digitising.

Snowline elevation The snowline elevation is synonymous with ELA when measured at the end of summer. All other snowline elevations apply to a transient seasonal snowline..

Total Area The entire area of the glacier. This may change from year to year, especially on the smaller glaciers.

Appendix A Index Glacier ELAs (1977-2011)

GLACIER	GL.IN. No	ELAo	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988
KAIKOURA RA	621/001	2490						5		-30				-5
MT. ELLA	932B/012	2142						12						
MT FAERIE QUEENE	646/006	2030						-45						5
MT. WILSON	None	1912	-37	-11			15		-115	-141	-63	41	-89	68
MT. FRANKLIN	911A/002	1814		-12	122	-41	-74		-148			-18	-54	35
ROLLESTON GL.	911A/004	1763	-8		-8	6	-3	10	-123	-8	-8	13	15	-1
MT. CARRINGTON	646C/027	1715		-45		-22	-87	25	-150	-115		-10	45	55
MT. AVOCA	685F/004	1965		-55				-15		-95	-55	-5		10
MARMADUKE GL.	664C/012	1830	-43	-35	78	2	-28	30	-136	-122	-50	9	-5	65
RETREAT GL	906A/004	1742		-4		38	-50	14	-252		-4	20		
BROWNING RA	906A/001	1598		7		10	-34	7	-113		-2	17		
DOUGLAS GL	685B/001	2040		91		-163		-18	-214	-240	-223	-128		91
MT. BUTLER	685C/060	1840	-55	112	55	-34	-40	28	-200	-65	-70	10	-9	42
DAINTY GL	897/019	1954	-32	36		-7	-77	93	-99		-81	-73	-84	-12
KEA GL	897/007	1820		65		44	-85	134	-230		-120	-82	-83	-58
JASPUR GL	897/003	1725		43		63		-15	-155		-105	-42		-42
SIEGE GL	893A/006	1736	-64	-24		-76	-70	-46	-268		-94	35		-70
VERTEBRAE #12	893A/012	1864		7		36	-51	-14	-64		-56	-44		7
VERTEBRAE #25	893A/025	1840		30		25	-39	0	-62		-50	-27		3
RIDGE GL.	711L/024	2226		79			2	10	-15	-141	-32	-9		59
LANGDALE GL.	711/035	2186	34	89			79	119	-1	-226	-221	14	-1	4
TASMAN GL.	711/012	1790	-10	85	-90	20	-35	-30	-80	-95	-90	-10	-29	50
SALISBURY GL	888B/003	1810	17	17		32	-58	17	-92	-51	-76	-1	-35	-81
JALF GL	886/002	1790	-15	-10		-32	-65	5	-230	-78	-146	-20	-51	-32
CHANCELLOR DOME	882A/007	1756	96	95		77	-93	92	-211		-147	-78	-28	52
GLENMARY GL.	711F/006	2175		57		-95	-45	6	-40	-155	-67			5
BLAIR GL.	711D/038	1938		74		-75	-13	-51	-126	-80	-85		-68	17
MT MCKENZIE	711D/021	1904		46		6	-16	-2	-184	-62	-124	13	-14	
JACKSON GL.	868B/094	2070		28		-20		8	-38	-80	-56		-9	
JACK GL.	875/015	1907		31		23	-22	44	-157	-79	-32	28	-2	-32
MT. ST. MARY	711B/039	1926								-171		-91		-19
THURNEYSON GL	711B/012	1970	-40			-44	-27		-105	-88	-65	-32	-52	-20
BREWSTER GL.	868C/020	1935		25		-89	-80	36	-141	-135	-139	-93	-107	
MT. STUART	752I/104	1673	-86	57		-23	-67	3	-135		-53	-13	5	
LINDSAY GL	867/002	1730		8		-78	-49	51	-170		-64	38	-115	42
FOG PK	752E/051	1987				-71		35		-96		-57	-85	35
SNOWY CK	752C/103	2092		64		-68	66	-54	-59	-68	-72	-56	-67	-55
MT. CARIA	863B/001	1472		-30			-59	-48	-100		-49	53	-50	28
FINDLAY GL.	859/009	1693					-89	42	-111		-64	32	-71	-8
PARK PASS GL.	752B/048	1824		79		-16	-46	34	-62	-59	-122	39	19	
MT. LARKINS	752E/002	1945								-265		-53		105
BRYANT GL.	752B/025	1783	-43	101			-20	-3	-163	-163	-173	-13	-20	
AILSA MTS.	752B/013	1648		-5					-88	-53	-53	1	-23	
MT. GUNN	851B/057	1593	22	45		-64	-62	17	-115		-53		-38	
MT. GENDARME	797G/033	1616					-46	-43	-136		-94	59	34	
LLAWRENNY PKS.	846/035	1476		4			-68	-4	-132		-36		-22	
BARRIER PK.	797F/004	1596		116		-51	-73	-31	-218		-72		-41	
MT. IRENE	797D/001	1563		137					-156		-37		-37	
MERRIE RA.	797B/010	1515		140										
CAROLINE PK.	803/001	1380												
NUMBER		50	15	40	5	32	36	41	41	27	40	38	33	33
MEAN		1836	-18	39	31	-21	-42	11	-132	-110	-80	-14	-35	10
STD. DEV.		215	46	50	83	53	39	43	64	63	51	45	40	46
No. below ELA (+ve balance)			9	6	1	19	29	17	40	25	32	19	19	19
% with +ve M.B.			60	15	20	59	81	41	98	93	80	50	58	58

Shaded columns indicate years of positive mass balance

GLACIER	GL.IN. No	ELAo	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
KAIKOURA RA	621/001	2490	50	168			-57	-30	-60			30	15
MT. ELLA	932B/012	2142	28				-87	-72	-142	-15		17	108
MT FAERIE QUEENE	646/006	2030	5				-62	-57	-65	-57		-50	145
MT. WILSON	None	1912	93				-167	-102	-154		-149	41	113
MT. FRANKLIN	911A/002	1814	52				-164	-109	-156	-82		122	164
ROLLESTON GL.	911A/004	1763	7				-143	-18	-143	-13	-123	40	87
MT. CARRINGTON	646C/027	1715	5				-170	-128	-158	-125	-137	-39	190
MT. AVOCA	685F/004	1965	6				-115	-25	-45	12	-75	70	115
MARMADUKE GL.	664C/012	1830	-13				-175	-75	-155	-82	-129	143	164
RETREAT GL	906A/004	1742	36				-73	-120	-277	16	-97	47	146
BROWNING RA	906A/001	1598	14				-116	-71	-118	-28	-108	30	
DOUGLAS GL	685B/001	2040	172			-197	-260	-154	-231	-150	-223	220	240
MT. BUTLER	685C/060	1840	-34			-95	-136	-28	-190	-60	-176	68	98
DAINTY GL	897/019	1954	-32			-103	-176	-73	-111	-12	-92	98	176
KEA GL	897/007	1820	-36				-230	-92	-250	80	-152	78	200
JASPUR GL	897/003	1725	-29				-150	-120	-145	-90	-100	43	
SIEGE GL	893A/006	1736	-64			-323	-386	-116	-396	-72	-203	214	414
VERTEBRAE #12	893A/012	1864	-39	310		-96	-86	-57	-86	-57	-73	-9	226
VERTEBRAE #25	893A/025	1840	-20			-94	-84	-54	-84	-54	-71	-5	125
RIDGE GL.	711L/024	2226	51				-126	-56	-94	-88	-136	9	109
LANGDALE GL.	711I/035	2186	69			-226	-231	14	-226	-1	-226	142	134
TASMAN GL.	711I/012	1790	-30		-35	-100	-108	-20	-124	-35	-102	63	186
SALISBURY GL	888B/003	1810	-66			-129	-100	-38	-165	-58	-84	42	220
JALF GL	886/002	1790	-31			-210	-220	-38	-240	-8	-191	-3	260
CHANCELLOR DOME	882A/007	1756	-78				-206	-147	-211	-36	-176	92	209
GLENMARY GL.	711F/006	2175	5			-15	-68	-30	-35	-45	-130	20	115
BLAIR GL.	711D/038	1938	-62				15	-73	-85	2		34	152
MT MCKENZIE	711D/021	1904	8				-122	8	-99	6	-189	31	174
JACKSON GL.	868B/094	2070	10				-78	-25	-52	2	-54	12	95
JACK GL.	875/015	1907	-9			-90	-142	-27	-152	51	-102	33	101
MT. ST. MARY	711B/039	1926	46			-76	-141	-76	-156	-71	-84	-37	199
THURNEYSON GL	711B/012	1970	0			-40	-60	-66	-102	-32	-70	-5	142
BREWSTER GL.	868C/020	1935	-17	310		-84	-185	-145	-158	27	-156	47	345
MT. STUART	752I/104	1673	-10				-138	-33	-158	39	-106	-17	132
LINDSAY GL	867/002	1730	34				-175	-120	-180	45	-85	70	145
FOG PK	752E/051	1987	45				-93	-87	-99	-93	-97	111	135
SNOWY CK	752C/103	2092	11				-34	-58	-62	-35	-72	66	148
MT. CARIA	863B/001	1472	-43				-97	-72	-106	-82	-77	-55	178
FINDLAY GL.	859/009	1693	-51				-118	-59	-132	-61	-113	87	197
PARK PASS GL.	752B/048	1824	-30					-41	-189	-16	-79	56	131
MT. LARKINS	752E/002	1945	91				-275	-95	-315	-163	-312	105	270
BRYANT GL.	752B/025	1783	-30				-108	-55	-153	-103	-113	-5	227
AILSA MTS.	752B/013	1648	-36				-84	-52	-93	-64	-55	-27	182
MT. GUNN	851B/057	1593	-59				-108	-64	-122	-86	-73	-34	209
MT. GENDARME	797G/033	1616	-36				-114	-64	-198	-100	-126	32	188
LLAWRENNY PKS.	846/035	1476	-47				-116	-68	-176	-15	-155	2	194
BARRIER PK.	797F/004	1596	-71				-168	-118	-236	-126	-132	86	304
MT. IRENE	797D/001	1563	-26				-156	-51	-163	-65	-103	49	95
MERRIE RA.	797B/010	1515	30				-135	-95	-165	-90	-130	-70	173
CAROLINE PK.	803/001	1380					-160	1	-150	-78	-130	2	182
NUMBER		50	49	2	1	15	49	50	50	48	45	50	48
MEAN		1836	-3	239	-65	-125	-136	-67	-151	-45	-124	42	172
STD. DEV.		215	49	100		80	68	40	70	53	52	62	70
No. below ELA (+ve balance)			26	0	1	15	48	47	50	38	45	13	0
% with +ve M.B.			53	0	100	100	98	94	100	79	100	26	0

GLACIER	GL.IN. No	ELAo	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
KAIKOURA RA	621/001	2490	10	-15	25			-25		14	50	5	13	30
MT. ELLA	932B/012	2142	60	-22	46	-64		-77	17	27	38	30	38	
MT FAERIE QUEENE	646/006	2030	160	-55	55	-54		-60	-48	-33	55	45	-47	
MT. WILSON	None	1912	108	13	93	-121	-92	-82	93	40	93	88	11	98
MT. FRANKLIN	911A/002	1814	136	-104	121	-70		-106	126	-23	126	124	106	166
ROLLESTON GL.	911A/004	1763	105	-8	97	-94	-1	-23	37	32	52	51	5	102
MT. CARRINGTON	646C/027	1715	237	-135	210	-84	-120	-137	135	-45	155	155	-90	245
MT. AVOCA	685F/004	1965	65	-52	88	-52	-20	-53	35	3	85	10	5	105
MARMADUKE GL.	664C/012	1830	153	-57	140	-117	-27	-130	145	13	160	140	-20	164
RETREAT GL	906A/004	1742	133	-60	93	-78	18	-132	48	-22	68	63	18	
BROWNING RA	906A/001	1598		-30	62	-77	-28	-110	12	21	27	22	2	
DOUGLAS GL	685B/001	2040	250	-40	245	-134	-35	-163	240	-44	340	243	245	330
MT. BUTLER	685C/060	1840	148	-95	90	-147	-80	-176	75	-2	120	85	-33	215
DAINTY GL	897/019	1954	74	-58	126	-67	-44	-94	-9	-12	116	41	-34	216
KEA GL	897/007	1820	190	-150	195	-159	-90	-155	78	-36	150	200	-83	
JASPUR GL	897/003	1725		-95	195	-74	-95	-125	35	-41		195	40	
SIEGE GL	893A/006	1736	279	-126	239	-160	-116	-203	-72	-18	394	224	-72	444
VERTEBRAE #12	893A/012	1864	121	-59	131	-48	-57	-74	-34	-8	166	60	-24	226
VERTEBRAE #25	893A/025	1840	70	-33	80	-51	-45	-75	-6	-16	110	19	-6	155
RIDGE GL.	711L/024	2226	79	-116	74	-63	2	-61	84	30	99	79	54	
LANGDALE GL.	711I/035	2186	129	-226	99	-226	-1	-221	109	74	394	129	76	
TASMAN GL.	711I/012	1790	110	-80	105	-80	-40	-40	60	6	235	175	-20	320
SALISBURY GL	888B/003	1810	172	-95	50	-95	-78	-95	40	0	140	140	-30	285
JALF GL	886/002	1790	265	-85		-190	-170	-190	-5	2	260	250	-35	370
CHANCELLOR DOME	882A/007	1756	204	-186	104	-186	-151	-181	94	43	109	109	-6	244
GLENMARY GL.	711F/006	2175	70	-37	35	-60	-30	-65	30	15	105	35	5	130
BLAIR GL.	711D/038	1938	147	-96	67	-93	-88	-88	22	34	132	34	15	222
MT MCKENZIE	711D/021	1904	148	-99	56	-134	-122	-132	6	15	111	31	-2	181
JACKSON GL.	868B/094	2070	63	-54	33	-55	-54	-56	15	14	93	12	2	160
JACK GL.	875/015	1907	85	-109	78	-147		-112	28	16	83	73	-9	293
MT. ST. MARY	711B/039	1926	204	-146	189	-131	-71	-126	-31	-16	194	74	-41	254
THURNEYSON GL	711B/012	1970	162	-92	135	-97	-35	-70	10	-7	140	130	5	180
BREWSTER GL.	868C/020	1935	220	-165	115	-141	-155	-155	-65	-120	335	55	-5	350
MT. STUART	752I/104	1673	177	-83	142	-103	-108	-108	52	-66	127	117	4	185
LINDSAY GL	867/002	1730	140	-90	142	-123	-122	-118	50	-30	130	80	-20	148
FOG PK	752E/051	1987	121	-92	125	-25	-95	-97	45	13	118	45	45	135
SNOWY CK	752C/103	2092	68	-88	28	-51	-58	-72	68	-2	66	64	64	
MT. CARIA	863B/001	1472	188	-52	153	-50	-92	-98	43	-17	48	48	-74	
FINDLAY GL.	859/009	1693	152	-73	132	-81	-115	-113	-3	-32	139	112	-61	177
PARK PASS GL.	752B/048	1824	111	-76	86	-163	-159	-154	76	26	86	86	56	181
MT. LARKINS	752E/002	1945	255	-280	150	-205	-255	-285	125	145	245	244	235	270
BRYANT GL.	752B/025	1783	182	-118	87	-117	-108	-118	17	-8	167	-8	-28	267
AILSA MTS.	752B/013	1648	137	-55	37	-65	-58	-68	-3	-35	22	27	-33	132
MT. GUNN	851B/057	1593	217	-78	42	-68	-108	-108	17	-43	52	197	-78	227
MT. GENDARME	797G/033	1616	159	-131	34	-96	-133	-129	32	-40	33	34	-46	184
LLAWRENNY PKS.	846/035	1476	181	-71	137	-100	-96	-76	9	-37	104	124	-68	124
BARRIER PK.	797I/004	1596	207	-108	194	-148	-131	-131	104	-17	119	194	-41	259
MT. IRENE	797D/001	1563	100	-162	170	-146	-130	-160	70	-85	50	68	-90	170
MERRIE RA.	797B/010	1515		-103	175	-100	-110	-105	150	-56	155	160	-75	
CAROLINE PK.	803/001	1380		-105	195	-89	-110	-125	30	-44	50	40	0	170
NUMBER		50	46	50	49	49	45	50	49	50	49	50	50	39
MEAN		1836	147	-91	112	-104	-85	-113	45	-7	132	95	-3	208
STD. DEV.		215	62	54	58	46	54	51	59	41	89	70	67	83.96
No. below ELA (+ve balance)			0	49	0	49	43	50	10	29	0	1	28	0
% with +ve M.B.			0	98	0	100	96	100	20	58	0	2	56	0

Appendix B Index Glacier details

Rock Glacier



GLACIER DATA		
AREA	45.42	ha
Debris area	41.66	ha
Max Elev	2640	m
Min Elev	2200	m
Mean Elev	2420	m
Length	1.4	km
Elev Range	440	m
Gradient	0.31	

SNOWLINE DATA		
Aspect	S	
ELAo	2490	m
Max SL	2540	m, 1989
Min SL	2430	m, 1995
Mean SL	2491	m
SL Range	110	m
No. surveys	13	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	AREAS ACCUM.	AREAS DEBRIS	ACCUM. TOTAL	MASS AREA RATIO (AAR)	MASS BALANCE INDEX
	m	m	ha	ha	ha		
ELAo	2490	0	3.88	41.66	45.54	0.09	0
1977							
1978							
1979							
1980							
1981							
1982	2495	5	3.90	41.64	45.54	0.09	-5
1983							
1984	2460	-30	6.74	38.80	45.54	0.15	30
1985							
1986							
1987							
1988	2485	-5	5.14	40.40	45.54	0.11	5
1989	2540	50	0.90	44.64	45.54	0.02	-50
1990							
1991							
1992							
1993	2433	-57	12.84	32.70	45.54	0.28	57
1994	2460	-30	6.74	38.80	45.54	0.15	30
1995	2430	-60	13.29	32.25	45.54	0.29	60
1996					no visit		
1997					cloud		
1998	2520	30	2.29	43.25	45.54	0.05	-30
1999	2505	15	2.74	42.80	45.54	0.06	-15
2000	2500	10	3.44	42.10	45.54	0.08	-10
2001	2475	-15	6.52	38.90	45.42	0.14	15
2002	2515	25	2.32	43.10	45.42	0.05	-25
2003					No visit	cloud	
2004					No visit	cloud	
2005	2465	-25	6.74	38.80	45.54	0.15	25
2006					no visit		
2007	2504	14	2.84	42.70	45.54	0.06	-14
2008	2540	50	0.90	44.64	45.54	0.02	-50
2009	2495	5	3.90	41.64	45.54	0.09	-5
2010	2503	13	3.34	42.20	45.54	0.07	-13
2011	2520	30	2.29	43.25	45.54	0.05	-30
MEAN	2491	1.39	4.83	40.70	45.53	0.11	-1



Photograph 1: Kaikoura Range 13 March 2011, resolution reduced to 200dpi.



Glacierette

GLACIER DATA		
Aspect	E	
AREA	5.32	ha
Max Elev	2250	m
Min Elev	2080	m
Mean Elev	2165	m
Length	0.34	km
Elev Range	170	m
Gradient	0.5	

SNOWLINE DATA		
ELAo	2142	m
Max SL	2250	m, 2000
Min SL	2000	m, 1995
Mean SL	2139	m
SL Range	250	m
No. surveys	17	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	2142	0	3.19	1.45	5.32	0.60	0
1977							
1978							
1979							
1980							
1981							
1982	2154	12	5.32	0.00	5.32	1.00	-12
1983							
1984							
1985							
1986							
1987							
1988							
1989	2170	28	2.17	3.15	5.32	0.41	-28
1990							
1991							
1992							
1993	2055	-87	5.91	0.00	5.91	1.11	87
1994	2070	-72	5.74	0.00	5.74	1.08	72
1995	2000	-142	15.10	0.00	15.10	2.84	142
1996	2127	-15	4.42	0.90	5.32	0.83	15
1997					In cloud		
1998	2159	17	2.92	2.40	5.32	0.55	-17
1999	2250	108	0	5.17	5.17	0.00	-108
2000	2202	60	0.1	2.46	2.56	0.04	-60
2001	2120	-22	4.57	0.75	5.32	0.86	22
2002	2188	46	0.33	1.61	1.94	0.17	-46
2003	2078	-64	5.32	0.00	5.32	1.00	64
2004					No visit	cloud	
2005	2065	-77	5.80	0.00	5.80	1.00	77
2006	2159	17	2.92	2.40	5.32	0.55	-17
2007	2169	27	2.20	3.10	5.30	0.42	-23
2008	2180	38	0.74	2.20	2.94	0.25	-23
2009	2172	30	1.14	1.80	2.94	0.39	-23
2010	2180	38	2.44	0.50	2.94	0.83	-23
2011	>2250		0.00	1.20	1.20	0.00	-23
MEAN	2139	-3	3.53	1.45		0.73	7



Photograph 2: Mt Ella 13 March 2011, resolution reduced to 200dpi.



Glacierette

GLACIER DATA		
Aspect	SE	
AREA	5.74	ha
Max Elev	2200	m
Min Elev	1940	m
Mean Elev	2070	m
Length	0.36	km
Elev Range	260	m
Gradient	0.72	

SNOWLINE DATA		
ELAo	2030	m
Max SL	2145	m, 2000
Min SL	1920	m, 1995
Mean SL	1976	m
SL Range	225	m
No. Surveys	14	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE	DEPARTURE	AREAS	ACCUM.	MASS
	ELEVATION	FROM ELAo	ABL.	TOTAL	AREA RATIO
	m	m	ha	ha	(AAR)
ELA	2030	45	3.44	2.30	5.74
					0.60
1977					
1978					
1979					
1980					
1981					
1982	1940	-45	5.74	0.00	5.74
1983					1.00
1984					
1985					
1986					
1987					
1988	1990	5	4.55	1.19	5.74
1989	1990	5	4.55	1.19	0.79
1990					0.79
1991					
1992					
1993	1923	-62	6.09	0.00	6.09
1994	1928	-57	5.99	0.00	5.99
1995	1920	-65	8.04	0.00	8.04
1996	1928	-57	5.99	0.00	5.99
1997					1.00
1998	1935	-50	5.84	0.00	5.84
1999	2130	145	0.00	5.74	3.80
2000	2145	160	0.00	2.90	2.90
2001	1930	-55	4.57	0.00	4.57
2002	2040	55	3.09	2.65	5.74
2003	1931	-54	4.57	0.00	4.57
2004				no visit	cloud
2005	1925	-60	6.09	0.00	6.09
2006	1937	-48	5.74	0.00	5.74
2007	1952	-33	5.44	0.30	5.74
2008	2040	55	0.60	2.80	3.40
2009	2030	45	1.10	2.30	3.40
2010	1938	-47	3.46	0.00	3.46
2011	>2200		0.00	2.72	2.72
MEAN	1976	-9	4.29	1.09	0.78
					9



Photograph 3: Faerie Queen 13 March 2011, resolution reduced to 200dpi.

Not numbered

Mt WILSON

NZMS 260 sheet K33



Snow patch

GLACIER DATA		
Aspect	S	
AREA	17.58 to 0.68	ha
Max Elev	2030	m
Min Elev	1740	m
Mean Elev	1885	m
Length	N/A	
Elev Range	290	m
Gradient		

SNOWLINE DATA		
ELA _o	1912	@ 7.03 ha
Max SL	2025	m, 1999
Min SL	1745	m, 1993
Mean SL	1901	m
SL Range	280	m
No. Surveys	26	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELA _o	AREAS ACCUM.	ABL.	TOTAL	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
	m	m	ha	N/A	ha		
ELA	1912		10.55	N/A	7.03	N/A	
1977	1875	-37	6.28		6.28		37
1978	1901	-11	4.68		4.68		11
1979							
1980							
1981	1927	15	3.28		3.28		-15
1982							
1983	1797	-115	4.88		4.88		115
1984	1771	-141	14.78		14.78		141
1985	1849	-63	8.38		8.38		63
1986	1953	41	2.28		2.28		-41
1987	1823	-89	10.48		10.48		89
1988	1980	68	1.48		1.48		-68
1989	2005	93	0.70		0.70		-93
1990							
1991							
1992							
1993	1745	-167	17.13		17.13		167
1994	1810	-102	11.38		11.38		102
1995	1758	-154	15.98		15.98		154
1996							
1997	1763	-149	15.48		15.48		149
1998	1953	41	2.28		2.28		-41
1999	2025	113	0.13		0.13		-113
2000	2020	108	0.68		0.68		-108
2001	1925	13	3.56		3.56		-13
2002	2005	93	0.63		0.63		-93
2003	1791	-121	4.40		4.40		121
2004	1820	-92	10.55		10.55		92
2005	1830	-82	10.48		10.48		82
2006	2005	93	0.65		0.65		-93
2007	1952	40	2.25		2.25		-40
2008	2005	93	0.65		0.65		-93
2009	2000	88	0.88		0.88		-88
2010	1923	11	3.45		3.45		-11
2011	2010	98	0.63		0.63		-98
Mean	1901	-11	5.66		5.66		11



Photograph 4: Mt Wilson 12 March 2011, resolution reduced to 200dpi.



Cirque

GLACIER DATA		
Aspect	E	
AREA	8.85	ha
Max Elev	2010	m
Min Elev	1680	m
Mean Elev	1845	m
Length	0.5	km
Elev Range	330	m
Gradient	0.66	

SNOWLINE DATA		
ELAo	1814	m
Max SL	1978	m, 1999
Min SL	1650	m, 1993
Mean SL	1823	m
SL Range	328	m
No. Surveys	24	

MEASUREMENTS

YEAR	SNOWLINE	DEPARTURE		AREAS		ACCUM.	MASS
	ELEVATION	FROM ELAo	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	1814	0	7.45	1.40	8.85	0.84	0
1977							
1978	1802	-12	7.74	1.11	8.85	0.87	12
1979	1936	122	0.84	8.01	8.85	0.09	-122
1980	1773	-41	8.30	0.55	8.85	0.94	41
1981	1740	-74	8.60	0.25	8.85	0.97	74
1982							
1983	1666	-148	?	0.00			148
1984							
1985							
1986	1796	-18	8.00	0.85	8.85	0.90	18
1987	1760	-54	8.40	0.45	8.85	0.95	54
1988	1849	35	5.30	3.55	8.85	0.60	-35
1989	1866	52	4.62	4.23	8.85	0.52	-52
1990							
1991							
1992							
1993	1650	-164	13.04	0.00	13.04	1.47	164
1994	1705	-109	8.80	0.05	8.85	0.99	109
1995	1658	-156	12.00	0.00	12.00	1.38	156
1996	1732	-82	8.65	0.20	8.85	0.98	82
1997					cloud		
1998	1936	122	0.84	8.01	8.85	0.09	-122
1999	1978	164	0.26	8.59	8.20	0.03	-164
2000	1950	136	0.57	8.28	7.90	0.07	-136
2001	1710	-104	8.85	0.00	7.90	1.12	104
2002	1935	121	0.87	6.88	7.75	0.11	-121
2003	1744	-70	7.50	0.25	7.75	0.97	70
2004					cloud		
2005	1708	-106	7.75	0.00	7.75	1.00	106
2006	1940	126	0.85	6.90	7.75	0.11	-126
2007	1791	-23	7.75	0.00	7.75	1.00	23
2008	1940	126	0.85	6.90	7.75	0.11	-126
2009	1938	124	0.80	6.95	7.75	0.10	-124
2010	1920	106	6.50	1.22	7.72	0.84	-106
2011	1980	166	0.10	7.62	7.72	0.01	-166
Mean	1823	9	5.51	2.93		0.68	-9



Photograph 5: Mt Franklin 12 March 2011, resolution reduced to 200dpi.



Cirque glacier

GLACIER DATA		
Aspect	SE	
AREA	10.8	ha
Max Elev	1900	m
Min Elev	1710	m
Mean Elev	1805	m
Length	0.36	km
Elev Range	190	m
Gradient	0.53	

SNOWLINE DATA		
ELAo	1763	m
Max SL	1868	m, 1993, 95
Min SL	1620	m, 2000
Mean SL	1761	m
SL Range	248	m
No. Surveys	30	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE	DEPARTURE		AREAS		ACCUM.	MASS
	ELEVATION	FROM ELAo	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
1955	1800	29					
1960	1800	29					-29
ELAo	1763	0	8.09	2.65	10.74	0.75	0
1977	1755	-8	9.10	1.64	10.74	0.85	8
1978	1770	7	8.19	2.55	10.74	0.76	-7
1979	1755	-8	9.10	1.64	10.74		8
1980	1769	6	8.19	2.55	10.74	0.76	-6
1981	1760	-3	8.60	2.14	10.74	0.80	3
1982	1773	10	8.09	2.65	10.74	0.75	-10
1983	1640	-123	10.74	0.00	10.74	1.00	123
1984	1755	-8	9.10	1.64	10.74	0.85	8
1985	1755	-8	9.10	1.64	10.74	0.85	8
1986	1776	13	7.94	2.80	10.74	0.74	-13
1987	1778	15	7.84	2.90	10.74	0.73	-15
1988	1762	-1	8.54	2.20	10.74	0.80	1
1989	1770	7	8.19	2.55	10.74	0.76	-7
1990							
1991							
1992							
1993	1620	-143	10.74	0.00	10.74	1.00	143
1994	1745	-18	9.90	0.84	10.74	0.92	18
1995	1620	-143	10.8	0.00	10.74	1.01	143
1996	1750	-13	9.49	1.25	10.74	0.88	13
1997	1640	-123	10.74	0.00	10.74	1.00	123
1998	1803	40	4.98	5.76	10.74	0.46	-40
1999	1850	87	1.3	9.44	10.74	0.12	-87
2000	1868	105	0.41	10.33	10.74	0.04	-105
2001	1755	-8	9.10	1.64	10.74	0.85	8
2002	1860	97	0.61	10.13	10.74	0.06	-97
2003	1669	-94	10.74	0.00	10.74	1.00	94
2004	1762	-1	8.54	2.20	10.74	0.80	1
2005	1740	-23	10.14	0.60	10.74	0.94	23
2006	1800	37	6.00	4.74	10.74	0.56	-37
2007	1795	32	6.31	4.43	10.74	0.59	-32
2008	1815	52	4.32	6.42	10.74	0.40	-52
2009	1814	51	4.39	6.35	10.74	0.41	-51
2010	1768	5	8.24	2.50	10.74	0.77	-5
2011	1865	102	0.50	10.24	10.74	0.05	-102
MEAN	1761	-2	7.72	3.02		0.71	2




Photograph 6: Rolleston Glacier 12 March 2011, resolution reduced to 200dpi.

GLACIER DATA		
Aspect	S	
AREA	15.5	ha
Max Elev	1960	m
Min Elev	1595	m
Mean Elev	1778	m
Length	0.71	km
Elev Range	365	m
Gradient	0.51	

SL DATA		
ELAo	1715	m
Max SL	1952	m, 2000
Min SL	1545	m, 1993
Mean SL	1703	m
SL Range	407	m
No. Surveys	28	

MEASUREMENTS Digitised values shaded

MEASUREMENTS							
YEAR	SNOWLINE	DEPARTURE	AREAS		ACCUM.	MASS	
	ELEVATION	FROM ELAo	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	1715	0	7.00	8.5	15.5	0.45	0
1977							
1978	1670	-45	8.86	6.64	15.50	0.57	45
1979							
1980	1693	-22	7.70	7.80	15.50	0.50	22
1981	1628	-87	11.60	3.90	15.50	0.75	87
1982	1740	25	6.50	9.00	15.50	0.42	-25
1983	1565	-150	18.50	0.00	15.50	1.19	150
1984	1600	-115	14.50	1.00	15.50	0.94	115
1985							
1986	1705	-10	7.20	8.30	15.50	0.46	10
1987	1760	45	6.00	9.50	15.50	0.39	-45
1988	1770	55	5.80	9.70	15.50	0.37	-55
1989	1720	5	6.90	8.60	15.50	0.45	-5
1990							
1991							
1992							
1993	1545	-170	20.55	0.00	15.50	1.33	170
1994	1587	-128	15.70	0.00	15.50	1.01	128
1995	1557	-158	16.60	0.00	15.50	1.07	158
1996	1590	-125	14.90	0.60	15.50	0.96	125
1997	1578	-137	17.43	0.00	15.50	1.12	137
1998	1676	-39	8.69	6.81	15.50	0.56	39
1999	1905	190	2.94	12.56	15.50	0.19	-190
2000	1952	237	0.48	15.02	15.50	0.03	-237
2001	1580	-135	17.30	0.00	15.50	1.12	135
2002	1925	210	1.30	6.70	8.00	0.16	-210
2003	1631	-84	6.60	1.40	8.00	0.83	84
2004	1595	-120	8.00	0.00	8.00	1.00	120
2005	1578	-137	17.43	0.00	15.50	1.12	137
2006	1850	135	4.20	11.30	15.50	0.27	-135
2007	1670	-45	8.86	6.64	15.50	0.57	45
2008	1870	155	3.70	11.80	15.50	0.24	-155
2009	1870	155	3.70	11.80	15.50	0.24	-155
2010	1625	-90	12.20	3.30	15.50	0.79	90
2011	1960	245	0.10	7.40	7.50	0.01	-245
MEAN	1703	-12	9.79	5.44		0.67	12





Photograph 7: Carrington 12 March 2011, resolution reduced to 200dpi.

GLACIER DATA		
Aspect	E	
AREA	9.803	m
Max Elev	2080	m
Min Elev	1890	m
Mean Elev	1985	m
Length	0.36	km
Elev Range	190	m
Gradient	0.53	

SNOWLINE DATA		
ELAo	1965	m
Max SL	2080	m, 1999
Min SL	1850	m, 1993
Mean SL	1963	m
SL Range	230	m
No. Surveys	26	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELA	1965	0	4.40	5.40	9.80	0.45	0
1977							
1978	1910	-55	9.20	0.60	9.80	0.94	55
1979							
1980							
1981							
1982	1950	-15	5.88	3.92	9.80	0.60	15
1983							
1984	1870	-95	10.50	0.00	9.80	1.00	95
1985	1910	-55	9.20	0.60	9.80	0.94	55
1986	1960	-5	4.85	4.95	9.80	0.49	5
1987							
1988	1975	10					-10
1989	1971	6	3.97	5.83	9.80	0.40	-6
1990							
1991							
1992							
1993	1850	-115	11.38	0.00	11.38	1.16	115
1994	1940	-25	7.20	2.60	9.80	0.73	25
1995	1920	-45	8.80	1.00	9.80	0.90	45
1996	1977	12	3.57	6.23	9.80	0.36	-12
1997	1890	-75	9.80	0.00	9.80	1.00	75
1998	2035	70	0.98	5.33	6.31	0.16	-70
1999	2080	115	0.00	3.22	3.22	0.00	-115
2000	2030	65	1.11	2.47	3.58	0.31	-65
2001	1913	-52	0.00	7.13	7.13	0.00	52
2002	2053	88	0.44	3.14	3.58	0.12	-88
2003	1913	-52	0.00	7.13	7.13	0.00	52
2004	1945	-20	6.50	3.30	9.80	0.66	20
2005	1912	-53	0.00	7.13	7.13	0.00	53
2006	2000	35	1.78	1.80	3.58	0.50	-35.00
2007	1968	3	2.38	1.20	3.58	0.66	-3.00
2008	2050	85	0.45	3.13	3.58	0.13	-85.00
2009	1975	10	2.28	1.30	3.58	0.64	-10.00
2010	1970	5	2.31	1.27	3.58	0.65	-5.00
2011	2070	105	0.10	0.97	1.07	0.09	-105.00
MEAN	1963	-2	4.27	3.05		0.51	2



Photograph 8: Mt Avoca 12 March 2011, resolution reduced to 200dpi.




Mountain glacier

GLACIER DATA		
Aspect	E	
AREA	93	m
Max Elev	2100	m
Min Elev	1615	m
Mean Elev	1858	m
Length	1.7	km
Elev Range	485	m
Gradient	0.285	

SNOWLINE DATA		
ELAo	1830	m
Max SL	1998	m, 1990
Min SL	1655	m, 1993
Mean SL	1835	m
SL Range	343	m
No. Surveys	33	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	1830	0	64	29	93	0.69	0
1977	1788	-43	73.02	19.50	92.52	0.79	43
1978	1795	-35	71.52	21.00	92.52	0.77	35
1979	1908	78	44.50	48.02	92.52	0.48	-78
1980	1832	2	63.52	29.00	92.52	0.69	-2
1981	1803	-28	69.72	22.80	92.52	0.75	28
1982	1860	30	49.65	42.87	92.52	0.54	-30
1983	1694	-136	86.72	5.80	92.52	0.94	136
1984	1708	-122	85.52	7.00	92.52	0.92	122
1985	1780	-50	73.97	18.55	92.52	0.80	50
1986	1839	9	61.72	30.80	92.52	0.67	-9
1987	1825	-5	64.77	27.75	92.52	0.70	5
1988	1895	65	48.02	44.50	92.52	0.52	-65
1989	1818	-13	67.02	25.50	92.52	0.72	13
1990	1998	168	22.52	70.00	92.52	0.24	-168
1991							
1992							
1993	1655	-175	90.34	2.18	92.52	0.98	175
1994	1755	-75	78.48	14.04	92.52	0.85	75
1995	1675	-155	92.14	0.38	92.52	1.00	155
1996	1748	-82	82.22	10.30	92.52	0.89	82
1997	1701	-129	91.87	0.65	92.52	0.99	129
1998	1973	143	26.81	65.71	92.52	0.29	-143
1999	1994	164	21.50	71.02	92.52	0.23	-164
2000	1983	153	23.75	68.77	92.52	0.26	-153
2001	1773	-57	75.32	17.20	92.52	0.81	57
2002	1970	140	28.38	53.42	81.80	0.35	-140
2003	1713	-117	74.50	7.30	81.80	0.91	117
2004	1803	-27	66.48	19.37	85.85	0.77	27
2005	1700	-130	91.87	0.65	92.52	0.99	130
2006	1975	145	27.80	54.00	81.80	0.34	-145
2007	1843	13	54.00	27.80	81.80	0.66	-13
2008	1990	160	26.80	55.00	81.80	0.33	-160
2009	1970	140	28.38	53.42	81.80	0.35	-140
2010	1810	-20	61.00	20.80	81.80	0.75	20
2011	1994	164	21.50	60.30	81.80	0.26	-164
MEAN	1835	5	60.09	30.14		0.66	-5



Photograph 9: Marmaduke Dixon 12 March 2011, resolution reduced to 200dpi.



Mountain glacier

GLACIER DATA		
Aspect	SW	
AREA	28.7	ha
Max Elev	1930	m
Min Elev	1570	m
Mean Elev	1750	m
Length	1.05	km
Elev Range	360	m
Gradient	0.343	

SNOWLINE DATA		
ELAo	1742	m
Max SL	1888	m, 1999
Min SL	1465	m, 1995
Mean SL	1726	m
SL Range	423	m
No. Surveys	27	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE DEPARTURE	AREAS	ACCUM.	MASS
	ELEVATION FROM ELAo	ABL.	TOTAL	AREA RATIO
	m	m	ha	ha
ELAo	1742	0	13.90	14.80
1977				
1978	1738	-4	14.70	14.00
1979				
1980	1780	38	8.29	20.41
1981	1692	-50	20.30	8.40
1982	1756	14	13.30	15.40
1983	1490	-252	31.59	-2.89
1984				
1985	1738	-4	14.83	13.87
1986	1762	20	10.78	17.92
1987				
1988				
1989	1778	36	8.87	19.83
1990				
1991				
1992				
1993	1669	-73	22.82	5.88
1994	1622	-120	27.20	1.50
1995	1465	-277	32.41	-3.71
1996	1758	16	11.65	17.05
1997	1645	-97	25.20	3.50
1998	1789	47	7.28	21.42
1999	1888	146	1.91	26.79
2000	1875	133	2.39	5.57
2001	1682	-60	0.69	7.27
2002	1835	93	2.88	5.08
2003	1664	-78	7.16	0.80
2004	1760	18	2.16	5.80
2005	1610	-132	6.46	1.50
2006	1790	48	0.36	7.60
2007	1720	-22	5.06	2.90
2008	1810	68	0.64	7.32
2009	1805	63	0.90	6.60
2010	1760	18	1.70	5.80
2011	>1888		0.00	5.25
MEAN	1726	-16	10.83	8.92



Photograph 10: Retreat Glacier on 13 March 2011, resolution reduced to 200dpi.



Small cirque

GLACIER DATA		
Aspect	S	
AREA	4.08	ha
Max Elev	1700	m
Min Elev	1530	m
Mean Elev	1615	m
Length		km
Elev Range	170	m
Gradient		

SNOWLINE DATA		
ELAo	1598	m
Max SL	1660	m, 2000
Min SL	1480	m, 1995
Mean SL	1573	m
SL Range	180	m
No. Surveys	27	

MEASUREMENTS Digitised values in bold type

MEASUREMENTS							
YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM.	AREAS	TOTAL	ACCUM.	MASS
	m	m	ha	ABL.	ha	AREA RATIO (AAR)	BALANCE INDEX
ELAo	1598	0	1.01	3.07	4.08	0.25	0
1977							
1978	1605	7	0.78	3.3	4.08	0.19	-7
1979							
1980	1608	10	0.74	3.34	4.08	0.18	-10
1981	1564	-34	2.45	1.63	4.08	0.60	34
1982	1605	7	0.78	3.3	4.08	0.19	-7
1983	1485	-113	7.48	0	7.48	1.00	113
1984							
1985	1596	-2	1.01	3.07	4.08	0.25	2
1986	1615	17	0.68	3.4	4.08	0.17	-17
1987							
1988							
1989	1612	14	2.01	2.07	4.08	0.49	-14
1990							
1991							
1992							
1993	1482	-116	2.03	2.05	4.08	0.50	116
1994	1527	-71	4.33	0	4.33	1.06	71
1995	1480	-118	20.38	0	20.38	5.00	118
1996	1570	-28	1.38	2.7	4.08	0.34	28
1997	1490	-108	12.81	0	12.81	1.00	108
1998	1628	30	0.55	3.53	4.08	0.13	-30
1999	>>1790		0	above gl	2.27	0.00	
2000	>>1790		0	above gl	1.38	0.00	
2001	1568	-30	2.18	1.9	3.58	0.61	30
2002	1660	62	0	1.14	1.14	0.00	-62
2003	1521	-77	4.08	0	4.08	1.00	77
2004	1570	-28	3.58	0	3.58	1.00	28
2005	1488	-110	12.81	0	12.81	1.00	110
2006	1610	12	0.33	3.25	3.58	0.09	-12.00
2007	1619	21	0.08	3.5	3.58	0.02	-21.00
2008	1625	27	0.05	2.55	2.60	0.02	-27.00
2009	1620	22	0.05	2.45	2.50	0.02	-22.00
2010	1600	2	0.10	1.49	1.59	0.06	-2.00
2011	>1790		0.00	0.88	0.88	0.00	
MEAN	1573	-25	2.99	1.82		0.55	25





Photograph 11: Browning Range 13 March 2011, resolution reduced to 200dpi.

GLACIER DATA		
Aspect	SE	
AREA	33.93	ha
Max Elev	2440	m
Min Elev	1820	m
Mean Elev	2130	m
Length	1.18	km
Elev Range	620	m
Gradient	0.508	

SNOWLINE DATA		
ELAo	2040	m
Max SL	2380	m, 2000
Min SL	1780	m, 1993
Mean SL	2045	m
SL Range	600	m
No. Surveys	29	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM.	AREAS	TOTAL	ACCUM.	MASS
	m	m	ha	ha	ha	AREA RATIO (AAR)	BALANCE INDEX
1974	2450				14.04		
ELAo	2040	0	24.73	9.20	33.93	0.73	0
1977							
1978	2131	91	19.93	14.00	33.93	0.59	-91
1979							
1980	1877	-163	32.03	1.90	33.93	0.94	163
1981							
1982	2022	-18	25.80	8.13	33.93	0.76	18
1983	1826	-214	33.73	0.20	33.93	0.99	214
1984	1800	-240	35.73	-1.80	33.93	1.05	240
1985	1817	-223	34.23	-0.30	33.93	1.01	223
1986	1912	-128	30.13	3.80	33.93	0.89	128
1987							
1988	2131	91	19.93	14.00	33.93	0.59	-91
1989	2212	172	15.74	18.19	33.93	0.46	-172
1990							
1991							
1992	1843	-197	33.13	0.80	33.93	0.98	197
1993	1780	-260	42.42	-8.49	33.93	1.25	260
1994	1886	-154	31.23	2.70	33.93	0.92	154
1995	1809	-231	35.43	-1.50	33.93	1.04	231
1996	1890	-150	31.43	2.50	33.93	0.93	150
1997	1817	-223	34.23	-0.30	33.93	1.01	223
1998	2260	220	10.71	23.22	33.93	0.32	-220
1999	2280	240	8.91	25.02	33.93	0.26	-240
2000	2290	250	7.96	25.97	33.93	0.23	-250
2001	2000	-40	22.21	6.60	28.81	0.77	40
2002	2285	245	3.31	25.50	28.81	0.11	-245
2003	1906	-134	25.61	3.20	28.81	0.89	134
2004	2005	-35	22.20	7.20	28.81	0.77	35
2005	1877	-163	28.80	0.00	28.80	1.00	163
2006	2280	240	8.91	19.90	28.81	0.31	-240
2007	1996	-44	22.41	6.40	28.81	0.78	44
2008	2380	340	21.20	0.90	22.10	0.96	-340
2009	2283	243	3.30	18.80	22.10	0.15	-243
2010	2340	300	3.97	25.55	22.06	0.18	-300
2011	2370	330	1.83	20.23	22.06	0.08	-330
Mean	2045	5	22.29	8.65		0.70	7





Photograph 12: Douglas Glacier 13 March 2011, resolution reduced to 200dpi.

GLACIER DATA		
Aspect	E	
AREA	75	ha
Max Elev	2040	m
Min Elev	1680	m
Mean Elev	1860	m
Length	1	km
Elev Range	360	m
Gradient	0	

SL DATA		
ELAo	1840	m
Max SL	2055	
Min SL	1640	m, 1995
Mean SL	1823	m
SL Range	415	m
No. surveys	33	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	ACCUM. TOTAL ha	MASS AREA RATIO (AAR)	MASS BALANCE INDEX
	m	m	ha	ha	ha		
ELAo	1840	0	42.76	31.80	74.56	0.57	0
1977	1785	-55	60.69	13.87	74.56	0.81	55
1978	1952	112	6.39	68.17	74.56	0.09	-112
1979	1895	55	21.46	53.10	74.56	0.29	-55
1980	1806	-34	55.76	18.80	74.56	0.75	34
1981	1800	-40	57.44	17.12	74.56	0.77	40
1982	1868	28	32.10	42.46	74.56	0.43	-28
1983	1640	-200	89.56	0.00	89.56	1.00	200
1984	1775	-65	62.66	11.90	74.56	0.84	65
1985	1770	-70	63.61	10.95	74.56	0.85	70
1986	1850	10	39.36	35.20	74.56	0.53	-10
1987	1831	-9	46.76	27.80	74.56	0.63	9
1988	1882	42	26.56	48.00	74.56	0.36	-42
1989	1806	-34	55.76	18.80	74.56	0.75	34
1990							
1991							
1992	1745	-95	67.36	7.20	74.56	0.90	95
1993	1704	-136	72.06	2.50	74.56	0.97	136
1994	1812	-28	53.56	21.00	74.56	0.72	28
1995	1650	-190	94.10	0.00	74.56	1.26	190
1996	1780	-60	62.06	12.50	74.56	0.83	60
1997	1664	-176	76.56	0.00	76.56	1.00	176
1998	1908	68	16.05	58.51	74.56	0.22	-68
1999	1938	98	8.04	53.66	61.70	0.13	-98
2000	1988	148	3.58	58.12	61.70	0.06	-148
2001	1745	-95	54.44	7.26	61.70	0.88	95
2002	1930	90	8.72	52.98	61.70	0.14	-90
2003	1693	-147	60.50	1.20	61.70	0.98	147
2004	1760	-80	53.57	8.13	61.70	0.87	80
2005	1664	-176	61.70	0.00	61.70	1.00	176
2006	1915	75	14.70	47.00	61.70	0.24	-75
2007	1862	22	34.44	16.43	50.87	0.68	-22
2008	1960	120	13.87	37.00	50.87	0.27	-120
2009	1925	85	10.37	40.50	50.87	0.20	-85
2010	1807	-33	45.17	5.70	50.87	0.89	33
2011	2055	215	0.32	39.61	39.93	0.01	-215
Mean	1823	-17	44.66	24.87	68.04	0.64	17




Photograph 13: Mt Butler 13 March 2011, resolution reduced to 200dpi.

GL DATA		
Aspect	W	
AREA	45.29	ha
Max Elev	2330	m
Min Elev	1750	m
Mean Elev	2040	m
Length	1.45	km
Elev Range	580	m
Gradient	0.4	

SL DATA		
ELA	1954	m
Max SL	2170	m, 1999
Min SL	1778	m, 1993
Mean SL	1941	m
SL Range	392	m
No. Surveys	31	

MEASUREMENTS Digitised values in bold type

YEAR	SNOWLINE	DEPARTURE		AREAS		ACCUM.	MASS
	ELEVATION	FROM ELA	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
1955	2100						
ELAo	1954	0	18.29	27	45.29	0.40	0
1977	1922	-32	25.81	19.48	45.29	0.57	32
1978	1990	36	14.79	30.5	45.29	0.33	-57
1979							
1980	1947	-7	20.19	25.1	45.29	0.45	7
1981	1877	-77	37.59	7.7	45.29	0.83	77
1982	2047	93	10.49	34.8	45.29	0.23	-93
1983	1855	-99	40.89	4.4	45.29	0.90	99
1984							
1985	1873	-81	38.29	7	45.29	0.85	81
1986	1881	-73	36.92	8.37	45.29	0.82	73
1987	1870	-84	39.09	6.2	45.29	0.86	84
1988	1942	-12	20.96	24.33	45.29	0.46	12
1989	1922	-32	25.81	19.48	45.29	0.57	32
1990							
1991							
1992	1851	-103	41.39	3.9	45.29	0.91	103
1993	1778	-176	43.71	1.58	45.29	0.97	176
1994	1881	-73	36.92	8.37	45.29	0.82	73
1995	1843	-111	41.94	3.35	45.29	0.93	111
1996	1942	-12	20.96	24.33	45.29	0.46	12
1997	1862	-92	40.29	5	45.29	0.89	92
1998	2052	98	10.23	35.06	45.29	0.23	-98
1999	2130	176	4.15	41.14	45.29	0.09	-176
2000	2028	74	11.79	33.5	45.29	0.26	-74
2001	1896	-58	32.95	12.34	45.29	0.73	58
2002	2080	126	8.25	37.04	45.29	0.18	-126
2003	1887	-67	36.09	9.2	45.29	0.80	67
2004	1910	-44	29.05	16.24	45.29	0.64	44
2005	1860	-94	40.89	4.4	45.29	0.90	94
2006	1945	-9	20.2	25.09	45.29	0.45	9
2007	1942	-12	20.96	24.33	45.29	0.46	12
2008	2070	116	8.99	36.3	45.29	0.20	-116
2009	1995	41	13.55	31.74	45.29	0.30	-41
2010	1920	-34	26.29	19	45.29	0.58	34
2011	2170	216	3.29	42	45.29	0.07	-216
MEAN	1941	-13	25.89	19.40	45.29	0.57	12





Photograph 14: Dainty Glacier 30 March 2011, resolution reduced to 200dpi..

GL DATA		
Aspect	S	
AREA	76.94	ha
Max Elev	2030	m
Min Elev	1650	m
Mean Elev	1840	m
Length	0.95	km
Elev Range	380	m
Gradient	0.453	

SL DATA		
ELAo	1820	m
Max SL	2020	m, 1999
Min SL	1570	m, 1995
Mean SL	1796	m
SL RANGE	450	m
No. Surveys	29	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	AREAS ABL.	ACCUM. TOTAL	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
	m	m	ha	ha	ha	
ELAo	1820	0	43.62	23.5	67.12	0.65
1977						
1978	1885	65	34.25	42.69	76.94	0.45
1979					no flight	
1980	1864	44	39.34	37.6	76.94	0.51
1981	1735	-85	66.64	10.3	76.94	0.87
1982	1954	134	16.53	60.41	76.94	0.21
1983	1590	-230	96.44	19.5	115.94	0.83
1984						
1985	1700	-120	73.1	3.84	76.94	0.95
1986	1738	-82	66.54	10.4	76.94	0.86
1987	1737	-83	66.59	10.35	76.94	0.87
1988	1762	-58	60.94	16	76.94	0.79
1989	1784	-36	56.43	20.51	76.94	0.73
1990						
1991						
1992						
1993	1590	-230	94.83	17.89	112.72	0.84
1994	1728	-92	68.13	8.81	76.94	0.89
1995	1570	-250	100.75	23.81	124.56	0.81
1996	1900	80	31.26	45.68	76.94	0.41
1997	1668	-152	80.79	3.85	84.63	0.95
1998	1898	78	31.74	45.2	76.94	0.41
1999	2020	200	1.47	75.47	76.94	0.02
2000	2010	190	2.3	74.64	76.94	0.03
2001	1670	-150	76.94	0	76.94	1.00
2002	2015	195	0.92	66.2	67.12	0.01
2003	1661	-159	67.12	0	67.12	1.00
2004	1730	-90	58.32	8.8	67.12	0.87
2005	1665	-155	67.12	0	67.12	1.00
2006	1898	78	31.74	35.38	67.12	0.47
2007	1784	-36	46.62	20.5	67.12	0.69
2008	1970	150	5.62	61.5	67.12	0.08
2009	2020	200	1.47	65.65	67.12	0.02
2010	1737	-83	59.12	8	67.12	0.88
2011	>2030		0	67.12	67.12	0.00
MEAN	1796	-24	51.63	27.67		0.64
						24





Photograph 15: Kea Glacier 30 March 2011, resolution reduced to 200dpi.

GLACIER DATA		
Aspect	SW	
AREA	14.09	ha
Max Elev	1920	m
Min Elev	1600	m
Mean Elev	1760	m
Length		km
Elev Range	320	m
Gradient		

SNOWLINE DATA		
ELAo	1725	m
Max SL	1920	m, 2000
Min SL	1570	m, 1983
Mean SL	1674	m
SL Range	350	m
No. Surveys	27	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1725	0	9.39	4.7	14.09	0.67	0
1977							
1978	1768	43	6.49	7.6	14.09	0.46	-43
1979					no flight		0
1980	1788	63	4.29	9.8	14.09	0.30	-63
1981							0
1982	1710	-15	10.34	3.75	14.09	0.73	15
1983	1570	-155	14.09	0	14.09	1.00	155
1984							0
1985	1620	-105	14.09	0	14.09	1.00	105
1986	1683	-42	11.84	2.25	14.09	0.84	42
1987							0
1988	1683	-42	11.84	2.25	14.09	0.84	42
1989	1696	-29	11.23	2.86	14.09	0.80	29
1990					no flight		0
1991					no flight		0
1992							0
1993	1575	-150	25.59	0	25.59	1.00	150
1994	1605	-120	17.39	0	17.39	1.00	120
1995	1580	-145	19.24	0	19.24	1.00	145
1996	1635	-90	13.59	0	13.59	1.00	90
1997	1625	-100	16.55	0	16.55	1.00	100
1998	1768	43	5.77	8.32	14.09	0.41	-43
1999	>1950			14.09	14.09	0.00	0
2000	>1950			7.84	7.84	0.00	0
2001	1630	-95	13.89	0.2	14.09	0.99	95
2002	1920	195	0	7.84	7.84	0.00	-195
2003	1651	-74	13.59	0	13.59	1.00	74
2004	1630	-95	13.39	0.2	13.59	0.99	95
2005	1600	-125	17.39	0	17.39	1.00	125
2006	1760	35	6.39	7.7	14.09	0.45	-35
2007	1684	-41	11.79	2.3	14.09	0.84	41
2008	> 1950		0	14.09	14.09	0.00	0
2009	1920	195	0	14.09	14.09	0.00	-195
2010	1765	40	6.09	8	14.09	0.43	-40
2011	> 1950		0	14.09	14.09	0.00	
MEANS	1690	-35	10.59	4.71	14.52	0.66	25



Photograph 16: Jaspur Group (eastern end only) 12 March 2011, resolution reduced to 200dpi.



GLACIER DATA		
Aspect	SE	
AREA	151.9	ha
Max Elev	2130	m
Min Elev	1370	m
Mean Elev	1750	m
Length	3.186	km
Elev Range	760	m
Gradient	0.239	

SNOW LINE DATA		
ELAo	1736	m
Max SL	2150	m, 1999
Min SL	1340	m, 1995
Mean SL	1675	m
SL Range	810	m
No. Surveys	30	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	AREAS ACCUM.	AREAS ABL.	ACCUM. TOTAL	MASS AREA RATIO (AAR)	MASS BALANCE INDEX
	m	m	ha	ha	ha		
ELAo	1736	0	61.4	90.5	151.9	0.40	0
1977	1672	-64	78.9	73	151.9	0.52	64
1978	1712	-24	67.66	84.24	151.9	0.45	24
1979					no flight		
1980	1660	-76	82.19	69.71	151.9	0.54	76
1981	1666	-70	80.4	71.5	151.9	0.53	70
1982	1690	-46	80.4	71.5	151.9	0.53	46
1983	1468	-268	130.57	21.33	151.9	0.86	268
1984							
1985	1642	-94	87.96	63.94	151.9	0.58	94
1986	1771	35	48.32	103.58	151.9	0.32	-35
1987							
1988	1666	-70	80.4	71.5	151.9	0.53	70
1989	1672	-64	78.81	73.09	151.9	0.52	64
1990					no flight		
1991					no flight		
1992	1413	-323	141.9	10	151.9	0.93	323
1993	1350	-386	153.9	0	153.9	1.00	386
1994	1620	-116	78.4	73.5	151.9	0.52	116
1995	1340	-396	155.4	0	155.4	1.00	396
1996	1664	-72	80.47	71.43	151.9	0.53	72
1997	1533	-203	116.4	35.5	151.9	0.77	203
1998	1950	214	30.83	121.07	151.9	0.20	-214
1999	2150	414	4.36	147.54	151.9	0.03	-414
2000	2015	279	24.27	127.63	151.9	0.16	-279
2001	1610	-126	97.61	54.29	151.9	0.64	126
2002	1975	239	28.9	123	151.9	0.19	-239
2003	1576	-160	107.4	44.5	151.9	0.71	160
2004	1620	-116	94.9	57	151.9	0.62	116
2005	1533	-203	116.4	35.5	151.9	0.77	203
2006	1664	-72	80.47	71.43	151.9	0.53	72
2007	1718	-18	66.2	85.7	151.9	0.44	18
2008	2130	394	5.8	146.1	151.9	0.04	-394
2009	1960	224	30.1	121.8	151.9	0.20	-224
2010	1664	-72	80.47	71.43	151.9	0.53	72
2011	2180	444	1.65	150.25	151.9	0.01	-444
MEANS	1709	-27	79.65	71.98		0.52	27



Photograph 17: Seige Glacier 30 March 2011, resolution reduced to 200dpi.



Cirque glacier

GLACIER DATA		
Aspect	SW	
AREA	20.64	ha
Max Elev	2100	m
Min Elev	1730	m
Mean Elev	1915	m
Length	0.63	km
Elev Range	370	m
Gradient	0.592	

SNOWLINE DATA		
ELAo	1864	m
Max SL	2090	m, 1999
Min SL	1768	m, 1992
Mean SL	1862	m
SL Range	322	m
No. surveys	29	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM.	AREAS	TOTAL	ACCUM.	MASS
	m	m	ha	ABL.	ha	AREA RATIO (AAR)	BALANCE INDEX
ELAo	1864	0	15.04	5.60	20.64	0.73	0
1977							
1978	1871	7	14.74	5.90	20.64	0.71	-7
1979					no flight		
1980	1900	36	12.94	7.70	20.64	0.63	-36
1981	1813	-51	17.54	3.10	20.64	0.85	51
1982	1850	-14	15.84	4.80	20.64	0.77	14
1983	1800	-64	18.27	2.37	20.64	0.89	64
1984							
1985	1808	-56	18.01	2.63	20.64	0.87	56
1986	1820	-44	17.34	3.30	20.64	0.84	44
1987					no visit		
1988	1871	7	14.74	5.90	20.64	0.71	-7
1989	1825	-39	26.96	3.49	20.64	1.31	39
1990					no flight		
1991					no flight		
1992	1768	-96	19.54	1.10	20.64	0.95	96
1993	1778	-86	19.14	1.50	20.64	0.93	86
1994	1807	-57	18.04	2.60	20.64	0.87	57
1995	1778	-86	19.13	1.51	20.64	0.93	86
1996	1807	-57	18.04	2.60	20.64	0.87	57
1997	1791	-73	18.64	2.00	20.64	0.90	73
1998	1855	-9	15.69	4.95	20.64	0.76	9
1999	2085	221	0.73	19.91	20.64	0.04	-221
2000	1985	121	5.53	15.11	20.64	0.27	-121
2001	1805	-59	18.06	2.58	20.64	0.88	59
2002	1995	131	4.94	15.70	20.64	0.24	-131
2003	1816	-48	17.49	3.15	20.64	0.85	48
2004	1807	-57	18.04	2.60	20.64	0.87	57
2005	1790	-74	18.66	1.98	20.64	0.90	74
2006	1830	-34	16.74	3.90	20.64	0.81	34
2007	1856	-8	15.54	5.10	20.64	0.75	8
2008	2030	166	3.14	17.50	20.64	0.15	-166
2009	1924	60	8.39	12.25	20.64	0.41	-60
2010	1840	-24	16.44	4.20	20.64	0.80	24
2011	2090	226	0.53	20.11	20.64	0.03	-226
MEAN	1862	-2	14.79	6.19	20.64	0.72	2



Photograph 18: Vertebrae Glaciers 12 30 March 2011, resolution reduced to 200dpi.

Mountain glacier

GLACIER DATA		
Aspect	SW	
Area	75.55	ha
Max Elev	2040	m
Min Elev	1700	m
Mean Elev	1850	m
Length	1.15 Km	km
Elev Range	360 m	m
Gradient	0.313	

SNOWLINE DATA		
ELAo	1840	m
Max SL	1995	m, 1999
Min SL	1746	m, 1992
Mean SL	1831	m
SL Range	249	m
No. surveys	29	


No 25

MEASUREMENTS Digitised values shaded

MEASUREMENTS

Digitized values shaded

YEAR	SNOWLINE	DEPARTURE		AREAS		ACCUM.	MASS
	ELEVATION	FROM ELA ₀	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELA ₀	1840	0	45.33	30.22	75.55	0.60	0
1977							
1978	1870	30	49.49	26.06	75.55	0.66	-30
1979							
1980	1865	25	52.12	23.43	75.55	0.69	-25
1981	1802	-39	70.44	5.11	75.55	0.93	39
1982	1840	0	65.53	10.02	75.55	0.87	0
1983	1778	-62	71.55	4.00	75.55	0.95	62
1984							
1985	1790	-50	70.55	5.00	75.55	0.93	50
1986	1813	-27	67.05	8.50	75.55	0.89	27
1987							
1988	1843	3	60.35	15.20	75.55	0.80	-3
1989	1820	-20	65.56	9.99	75.55	0.87	20
1990							
1991							
1992	1746	-94	73.65	1.90	75.55	0.97	94
1993	1756	-84	73.08	2.47	75.55	0.97	84
1994	1786	-54	71.05	4.50	75.55	0.94	54
1995	1756	-84	73.08	2.47	75.55	0.97	84
1996	1786	-54	71.05	4.50	75.55	0.94	54
1997	1770	-71	72.25	3.30	75.55	0.96	71
1998	1835	-5	61.98	13.57	75.55	0.82	5
1999	1965	125	2.64	72.91	75.55	0.03	-125
2000	1910	70	18.08	57.47	75.55	0.24	-70
2001	1807	-33	68.03	7.52	75.55	0.90	33
2002	1920	80	12.55	63.00	75.55	0.17	-80
2003	1789	-51	70.75	4.80	75.55	0.94	51
2004	1795	-45	70.55	5.00	75.55	0.93	45
2005	1765	-75	72.25	3.30	75.55	0.96	75
2006	1834	-6	61.98	13.57	75.55	0.82	6
2007	1824	-16	64.85	10.70	75.55	0.86	16
2008	1950	110	4.75	70.80	75.55	0.06	-110
2009	1859	19	36.72	38.83	75.55	0.49	-19
2010	1834	-6	61.98	13.57	75.55	0.82	6
2011	1995	155	0.64	74.91	75.55	0.01	-155
MEAN	1831	-9	55.67	19.88		0.74	15



Photograph 19: Vertebrae Glaciers 25 30 March 2011, resolution reduced to 200dpi.



Mountain glacier

GLACIER DATA		
Aspect	SE	
AREA	68.00	ha
Max Elev	2490	m
Min Elev	2110	m
Mean Elev	2300	m
Length	1.04	km
Elev Range	380	m
Gradient	0.365	

SNOWLINE DATA		
ELAo	2226	m
Max SL	2335	m,1999
Min SL	2085	m,1984
Mean SL	2222	m
SL Range	250	m
No. surveys	28	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE	DEPARTURE		AREAS		ACCUM.	MASS
	ELEVATION	FROM ELAo	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	2226	0	51.72	21.80	73.52	0.70	0
1977							
1978	2305	79	22.22	51.30	73.52	0.30	-79
1979					no flight		
1980							
1981	2228	2	49.41	24.11	73.52	0.67	-2
1982	2236	10	47.98	25.54	73.52	0.65	-10
1983	2211	-15	56.32	17.20	73.52	0.77	15
1984	2085	-141	73.52	0.00	73.52	1.00	141
1985	2194	-32	60.40	13.12	73.52	0.82	32
1986	2217	-9	55.02	18.50	73.52	0.75	9.1
1987							
1988	2285	59	27.82	45.70	73.52	0.38	-59
1989	2277	51	30.01	43.51	73.52	0.41	-51
1990					no flight		
1991					no flight		
1992							
1993	2100	-126	73.52	0.00	73.52	1.00	126
1994	2170	-56	66.32	7.20	73.52	0.90	56
1995	2132	-94	70.83	2.69	73.52	0.96	94
1996	2138	-88	70.01	3.51	73.52	0.95	88
1997	2090	-136	73.52	0.00	73.52	1.00	136
1998	2235	9	47.02	25.50	72.52	0.65	-9
1999	2335	109	19.92	52.00	71.92	0.28	-109
2000	2305	79	25.85	44.87	70.72	0.37	-79
2001	2110	-116	70.12	0.00	70.12	1.00	116
2002	2300	74	22.52	47.00	69.52	0.32	-74
2003	2163	-63	62.42	6.50	68.92	0.91	63
2004	2228	2	50.04	18.46	68.50	0.73	-2
2005	2165	-61	61.70	6.30	68.00	0.91	61
2006	2310	84	20.00	48.00	68.00	0.29	-84
2007	2256	30	38.30	29.70	68.00	0.56	-30
2008	2325	99	17.48	50.52	68.00	0.26	-99
2009	2305	79	34.11	44.87	68.00	0.50	-79
2010	2280	54	27.20	40.80	68.00	0.40	-54
2011	>2490		0.00	68.00	68.00	0.00	
MEAN	2222	-4	47.17	26.25		0.66	4



Photograph 20: Ridge Glacier 30 March 2011, resolution reduced to 200dpi.



Cirque glacier

GLACIER DATA		
Aspect	NW	
AREA	34.10	ha
Max Elev	2580	m
Min Elev	2090	m
Mean Elev	2335	m
Length	1.03	km
Elev Range	490	m
Gradient	0.478	

SNOWLINE DATA		
ELA	2186	m
Max SL	2328	m,1998
Min SL	1955	m,1984
Mean SL	2175	m
SL Range	373	m
No. surveys	29	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE	DEPARTURE	AREAS		ACCUM.	MASS	
	ELEVATION	FROM ELA	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	2186	0	29.42	11.00	40.42	0.73	0
1977	2220	34	32.62	7.80	40.42	0.81	-34
1978	2275	89	19.09	21.00	40.09	0.48	-89
1979					no flight		
1980							
1981	2265	79	15.12	24.31	39.43	0.38	-79
1982	2305	119	23.26	16.17	39.43	0.59	-119
1983	2185	-1	32.73	6.70	39.43	0.83	1
1984	1960	-226	39.43	0.00	39.43	1.00	226
1985	1965	-221	37.74	1.69	39.43	0.96	221
1986	2200	14	35.23	4.20	39.43	0.89	-14
1987	2185	-1	36.53	2.90	39.43	0.93	1
1988	2190	4	26.60	12.50	39.10	0.68	-4
1989	2255	69	18.37	20.73	39.10	0.47	-69
1990					no flight		
1991					no flight		
1992	1960	-226	38.11	0.00	38.11	1.00	226
1993	1955	-231	38.11	0.00	38.11	1.00	231
1994	2200	14	13.85	24.26	38.11	0.36	-14
1995	1960	-226	38.11	0.00	38.11	1.00	226
1996	2185	-1	32.11	6.00	38.11	0.84	1
1997	1960	-226	38.11	0.00	38.11	1.00	226
1998	2328	142	5.40	32.27	37.67	0.14	-142
1999	2320	134	6.68	30.00	36.68	0.18	-134
2000	2315	129	6.39	29.30	35.69	0.18	-129
2001	1960	-226	35.69	0.00	35.69	1.00	226
2002	2285	99	8.78	26.00	34.78	0.25	-99
2003	1960	-226	34.78	0.00	34.78	1.00	226
2004	2185	-1	26.00	8.78	34.78	0.75	1
2005	1965	-221	34.78	0.00	34.78	1.00	221
2006	2295	109	6.80	27.70	34.50	0.20	-109
2007	2260	74	14.50	20.00	34.50	0.42	-74
2008	2580	394	0.00	34.10	34.10	0.00	-394
2009	2315	129	6.39	27.71	34.10	0.19	-129
2010	2262	76	13.20	20.90	34.10	0.39	-76
2011	>2580		0.00	34.10	34.10	0.00	
MEAN	2175	-11	24	14.17		0.63	11





Photograph 21: Langdale Glacier 30 March 2011, resolution reduced to 200dpi.

GLACIER DATA		
Aspect	SW	
Max Elev		
Min Elev		
Mean Elev		
Elev Range		
Length		
Gradient		
Gradient @ S	0.096	

SNOWLINE DATA		
ELAo	1790	1955 to '01
ELA	1803	1977 to '01
Max SL	2110	m, 1990
Min SL	1666	m, 1995
Mean SL	1803	m
SL Range	444	m
No. surveys	35	

MEASUREMENTS Values from contour counts shaded

YEAR	SNOWLINE	DEPARTURE	AREAS		ACCUM.	MASS	
	ELEVATION	FROM ELA	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	1790	Too large to measure					
1977	1780	-10					10
1978	1875	85					-85
1979	1700	-90					90
1980	1810	20					-20
1981	1755	-35					35
1982	1760	-30					30
1983	1710	-80					80
1984	1695	-95					95
1985	1700	-90					90
1986	1780	-10					10
1987	1761	-29					29
1988	1840	50					-50
1989	1760	-30					30
1990	2100	310					-310
1991	1755	-35					35
1992	1690	-100					100
1993	1682	-108					108
1994	1770	-20					20
1995	1666	-124					124
1996	1755	-35					35
1997	1688	-102					102
1998	1853	63					-63
1999	1976	186					-186
2000	1900	110					-110
2001	1710	-80					80
2002	1895	105					-105
2003	1710	-80					80
2004	1750	-40					40
2005	1750	-40					40
2006	1850	60					-60
2007	1796	6					-6
2008	2025	235					-235
2009	1965	175					-175
2010	1770	-20					20
2011	2110	320					-320
MEAN	1803	13					-13



Photograph 22: Tasman Glacier 30 March 2011, resolution reduced to 200dpi.

Mountain glacier



GLACIER DATA		
Aspect	SW	
AREA	311.90	ha
Max Elev	2390	m
Min Elev	1340	m
Mean Elev	1865	m
Length	2.98	km
Elev Range	1050	m
Gradient	0.352	

SNOWLINE DATA		
ELAo	1810	m
Max SL	2095	m, 1999
Min SL	1645	m, 1995
Mean SL	1802	m
SL Range	450	m
No. surveys	32	

MEASUREMENTS



Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1810	0	246.90	65.00	311.90	0.79	0
1977	1827	17	239.90	72.00	311.90	0.77	-17
1978	1827	17	239.90	72.00	311.90	0.77	-17
1979					No flight		
1980	1842	32	228.51	83.39	311.90	0.73	-32
1981	1752	-58	268.90	43.00	311.90	0.86	58
1982	1827	17	239.90	72.00	311.90	0.77	-17
1983	1718	-92	279.56	32.34	311.90	0.90	92
1984	1760	-51	266.90	45.00	311.90	0.86	51
1985	1734	-76	275.90	36.00	311.90	0.88	76
1986	1809	-1	250.90	61.00	311.90	0.80	1
1987	1775	-35	261.90	50.00	311.90	0.84	35
1988	1729	-81	268.90	43.00	311.90	0.86	81
1989	1744	-66	271.90	40.00	311.90	0.87	66
1990					No flight		
1991					No flight		
1992	1681	-129	289.90	22.00	311.90	0.93	129
1993	1710	-100	279.90	32.00	311.90	0.90	100
1994	1772	-38	228.96	82.94	311.90	0.73	38
1995	1645	-165	296.67	15.23	311.90	0.95	165
1996	1752	-58	267.90	44.00	311.90	0.86	58
1997	1726	-84	276.90	35.00	311.90	0.89	84
1998	1852	42	223.24	88.66	311.90	0.72	-42
1999	2030	220	116.70	195.20	311.90	0.37	-220
2000	1982	172	132.50	179.40	311.90	0.42	-172
2001	1715	-95	281.66	30.24	311.90	0.90	95
2002	1860	50	219.00	92.00	311.00	0.70	-50
2003	1715	-95	279.90	32.00	311.90	0.90	95
2004	1732	-78	275.90	36.00	311.90	0.88	78
2005	1715	-95	281.66	30.24	311.90	0.90	95
2006	1850	40	223.24	88.66	311.90	0.72	-40
2007	1810	0	248.90	63.00	311.90	0.80	0
2008	1950	140	156.90	155.00	311.90	0.50	-140
2009	1950	140	156.90	155.00	311.90	0.50	-140
2010	1780	-30	259.90	52.00	311.90	0.83	30
2011	2095	285	61.84	250.06	311.90	0.20	-285
Mean	1802	-8	239	73		0.77	8





Photograph 23: Salisbury Snowfield 30 March 2011, resolution reduced to 200dpi.

No. 886/2 & 888B/7			JALF GL. and Baumann Gl.			NZMS 260 sheet H35	
			Small saddle glacier				
	GLACIER DATA (2008)				SNOWLINE DATA		
	Aspects	N &S, E & W					
	AREA	41.50	ha		ELAo	1790	m
	Max Elev	1880	m		Max SL	2160	m, 2000
	Min Elev	1600	m		Min SL	1550	m, 1995
	Mean Elev	1740	m		Mean SL	1762	m
	Length	na			SL Range	610	m
	Elev Range	280	m		No. surveys	31	
	Gradient	na					
Glacier area has reduced since the first map in 1977							
MEASUREMENTS		Digitised values shaded					
YEAR	SNOWLINE	DEPARTURE		AREAS		ACCUM.	MASS
	ELEVATION	FROM ELAo	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	1790	0	14.10	88.00	102.10	0.14	0
1977	1775	-15	21.30	80.80	102.10	0.21	15
1978	1780	-10	19.10	83.00	98.61	0.19	10
1979					95.12	no visit	
1980	1758	-32	36.41	58.71	95.12	0.38	32
1981	1725	-65	55.63	36.00	91.63	0.61	65
1982	1795	5	2.63	89.00	91.63	0.03	-5
1983	1560	-230	91.63	0.00	91.63	1.00	230
1984	1712	-78	72.96	18.67	91.63	0.80	78
1985	1644	-146	91.63	0.00	91.63	1.00	146
1986	1770	-20	13.93	77.70	91.63	0.15	20
1987	1739	-51	45.63	46.00	91.63	0.50	51
1988	1758	-32	22.14	66.00	88.14	0.25	32
1989	1759	-31	21.66	66.48	88.14	0.25	31
1990					no visit		
1991					no visit		
1992	1580	-210	77.67	0.00	77.67	1.00	210
1993	1570	-220	77.67	0.00	77.67	1.00	220
1994	1752	-38	17.04	60.63	77.67	0.22	38
1995	1550	-240	77.67	0.00	77.67	1.00	240
1996	1782	-8	17.25	60.42	77.67	0.22	8
1997	1599	-191	75.11	2.56	77.67	0.97	191
1998	1787	-3	15.82	57.20	73.02	0.22	3
1999	2050	260	0.00	102.10	62.55	0.00	-260
2000	2055	265	0.00	102.10	52.08	0.00	-265
2001	1705	-85	26.04	26.04	52.08	0.50	85
2002					43.00		
2003	1600	-190	43.00	3.00	43.00	1.00	190
2004	1620	-170	42.80	0.20	43.00	1.00	170
2005	1600	-190	40.00	3.00	43.00	0.93	190
2006	1785	-5	5.59	37.18	42.77	0.13	5.00
2007	1792	2	0.00	42.77	42.77	0.00	-2.00
2008	2050	260	0.00	41.50	41.50	0.00	-260.00
2009	2040	250	0.00	41.00	41.00	0.00	-250.00
2010	1755	-35	3.50	37.50	41.00	0.09	35.00
2011	2160	370	0.00	41.00	41.00	0.00	-370.00
MEAN	1762	-28	33	41.31		0.44	28



Photograph 24: Jalf Glacier 30 March 2011, resolution reduced to 200dpi.

No. 882A/7			CHANCELLOR DOME					NZMS 260 sheet H35 & H36
				Cirque glacier				
	GLACIER DATA				SNOWLINE DATA			
	Aspect	SW						
	AREA	17.51	ha		ELAo	1756	m	
	Max Elev	1960	m		Max SL	2000	m, 1999	
	Min Elev	1660	m		Min SL	1545	m, 1995	
	Mean Elev	1810	m		Mean SL	1739	m	
	Length	0.55	km		SL Range	455	m	
	Elev Range	300	m		No. surveys	30		
	Gradient	0.436						
MEASUREMENTS			Digitised values shaded					
YEAR	SNOWLINE	DEPARTURE		AREAS		ACCUM.	MASS	
	ELEVATION	FROM ELAo	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE	
	m	m	ha	ha	ha	(AAR)	INDEX	
ELAo	1756	0	18.83	5.25	24.08	0.78	0	
1977	1852	96	9.45	14.63	24.08	0.39	-96	
1978	1851	95	9.63	14.45	24.08	0.40	-95	
1979								
1980	1833	77	13.85	10.23	24.08	0.58	-77	
1981	1663	-93	21.05	3.03	24.08	0.87	93	
1982	1848	92	10.67	13.41	24.08	0.44	-92	
1983	1545	-211	28.90	0.00	28.90	1.20	211	
1984								
1985	1609	-147	22.95	1.13	24.08	0.95	147	
1986	1678	-78	20.65	3.43	24.08	0.86	78	
1987	1728	-28	19.33	4.75	24.08	0.80	28	
1988	1808	52	17.33	6.75	24.08	0.72	-52	
1989	1678	-78	20.59	3.49	24.08	0.86	78	
1990								
1991								
1992								
1993	1550	-206	24.08	0.00	24.08	1.00	206	
1994	1609	-147	22.95	1.13	24.08	0.95	147	
1995	1545	-211	28.90	0.00	28.90	1.00	211	
1996	1720	-36	7.42	16.66	24.08	0.31	36	
1997	1580	-176	24.08	0.00	24.08	1.00	176	
1998	1848	92	10.67	13.41	24.08	0.44	-92	
1999	1965	209	0.00	17.51	17.51	0.00	-209	
2000	1960	204	0.00	17.51	17.51	0.00	-204	
2001	1570	-186	17.51	0.00	17.51	1.00	186	
2002	1860	104	1.50	16.01	17.51	0.09	-104	
2003	1570	-186	17.51	0.00	17.51	1.00	186	
2004	1605	-151	17.51	0.00	17.51	1.00	151	
2005	1575	-181	17.51	0.00	17.51	1.00	181	
2006	1850	94	10.00	16.10	17.51	0.57	-94	
2007	1799	43	3.31	14.20	17.51	0.19	-43	
2008	1865	109	1.30	16.21	17.51	0.07	-109	
2009	1865	109	1.30	16.21	17.51	0.07	-109	
2010	1750	-6	10.51	7.00	17.51	0.60	6	
2011	2000	244	0.00	17.51	17.51	0.00	-244	
MEAN	1739	-17	14.76	7.56		0.63	17	



Photograph 25: Chancellor Glacier 30 March 2011, resolution reduced to 200dpi.



GLACIER DATA		
Aspect	S	
AREA	58.51	ha
Max Elev	2380	m
Min Elev	2040	m
Mean Elev	2210	m
Length	1.19	km
Elev Range	340	m
Gradient	0.277	

SNOWLINE DATA		
ELAo	2164	m
Max SL	2305	m, 1999
Min SL	2020	m, 1984
Mean SL	2165.206897	m
SL Range	285	m
No. Surveys	29	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	2175	0	35.41	19.00	54.41	0.65	0
1977							
1978	2232	57	22.93	31.48	54.41	0.42	-57
1979							
1980	2080	-95	48.50	5.57	54.07	0.90	95
1981	2130	-45	42.20	11.53	53.73	0.79	45
1982	2181	6	36.00	17.73	53.73	0.67	-6
1983	2135	-40	41.76	11.97	53.73	0.78	40
1984	2020	-155	53.73	0.00	53.73	1.00	155
1985	2108	-67	49.30	4.43	53.73	0.92	67
1986					53.73		
1987					53.73		
1988	2180	5	36.02	17.37	53.39	0.67	-5
1989	2180	5	36.02	17.37	53.39	0.67	-5
1990					52.39		
1991					52.39		
1992	2160	-15	37.80	14.59	52.39	0.72	15
1993	2107	-68	45.50	6.89	52.39	0.87	68
1994	2145	-30	40.00	12.39	52.39	0.76	30
1995	2140	-35	40.70	11.69	52.39	0.78	35
1996	2130	-45	42.25	10.14	52.39	0.81	45
1997	2045	-130	50.51	1.88	52.39	0.96	130
1998	2195	20	32.34	19.73	52.07	0.62	-20
1999	2290	115	7.70	43.92	51.62	0.15	-115
2000	2245	70	20.19	30.98	51.17	0.39	-70
2001	2138	-37	41.10	10.07	51.17	0.80	37
2002	2210	35	30.01	20.84	50.85	0.59	-35
2003	2115	-60	44.50	6.33	50.83	0.88	60
2004	2145	-30	40.00	10.81	50.81	0.79	30
2005	2110	-65	45.10	5.70	50.80	0.89	65
2006	2205	30	31.12	19.68	50.80	0.61	-30
2007	2190	15	32.40	18.40	50.80	0.64	-15
2008	2280	105	9.74	41.06	50.80	0.19	-105
2009	2210	35	29.96	20.84	50.80	0.59	-35
2010	2180	5	34.30	16.50	50.80	0.68	-5
2011	2305	130	5.26	45.54	50.80	0.10	-130
Mean	2165	-10	35.4	16.7		0.68	10





Photograph 26: Glenmary Glacier on 30 March 2011, resolution reduced to 200dpi.

GLACIER DATA		
Aspect	SE	
AREA	24.62	ha
Max Elev	2240	m
Min Elev	1790	m
Mean Elev	2015	m
Length	0.63	km
Elev Range	450	m
Gradient	0.71	

SNOWLINE DATA		
ELAo	1938	m
Max SL	2160	m, 1999
Min SL	1812	m, 1983
Mean SL	1934	m
SL Range	348	m
No. Surveys	28	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	ACCUM. TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1938	0	18.52	7.90	26.42	0.70	0
1977							
1978	2012	74	13.42	13.00	26.42	0.51	-74
1979							
1980	1863	-75	22.82	3.60	26.42	0.86	75
1981	1925	-13	19.23	7.19	26.42	0.73	13
1982	1887	-51	21.52	4.90	26.42	0.81	51
1983	1812	-126	25.28	1.14	26.42	0.96	126
1984	1858	-80	23.12	3.30	26.42	0.88	80
1985	1853	-85	23.28	3.14	26.42	0.88	85
1986							
1987	1870	-68	22.43	3.99	26.42	0.85	68
1988	1955	17	17.80	8.62	26.42	0.67	-17
1989	1876	-62	22.10	4.32	26.42	0.84	62
1990							
1991							
1992							
1993	1953	15	17.69	8.73	26.42	0.67	-15
1994	1865	-73	22.72	3.70	26.42	0.86	73
1995	1853	-85	23.28	3.14	26.42	0.88	85
1996	1940	2	18.44	7.98	26.42	0.70	-2
1997							
1998	1972	34	16.36	10.06	26.42	0.62	-34
1999	2090	152	6.04	20.38	26.42	0.23	-152
2000	2085	147	6.17	20.25	26.42	0.23	-147
2001	1842	-96	23.75	2.67	26.42	0.90	96
2002	2005	67	14.02	12.40	26.42	0.53	-67
2003	1845	-93	23.62	2.80	26.42	0.89	93
2004	1850	-88	23.28	3.14	26.42	0.88	88
2005	1850	-88	23.28	3.14	26.42	0.88	88
2006	1960	22	17.60	8.82	26.42	0.67	-22
2007	1972	34	16.36	10.06	26.42	0.62	-34
2008	2070	132	7.92	18.50	26.42	0.30	-132
2009	1972	34	16.36	10.06	26.42	0.62	-34
2010	1953	15	17.69	8.73	26.42	0.67	-15
2011	2160	222	2.52	23.90	26.42	0.10	-222
MEAN	1934	-4	18.15	7.69		0.69	4



Photograph 27: Blair Glacier on 30 March 2011, resolution reduced to 200dpi.

Mountain glacier



GLACIER DATA		
Aspect	S	
AREA	48.67	ha
Max Elev	2100	m
Min Elev	1760	m
Mean Elev	1930	m
Length	0.69	km
Elev Range	340	m
Gradient	0.49	

SNOWLINE DATA		
ELAo	1904	m
Max SL	2085	m, 1999
Min SL	1715	m, 1983
Mean SL	1888	m
SL Range	370	m
No. Surveys	29	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	ACCUM. TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1904	0	30.87	17.80	48.67	0.63	0
1977							
1978	1950	46	22.78	25.89	48.67	0.47	-46
1979							
1980	1910	6	29.86	18.81	48.67	0.61	-6
1981	1888	-16	33.48	15.19	48.67	0.69	16
1982	1902	-2	31.35	17.32	48.67	0.64	2
1983	1720	-184	60.63	-11.96	48.67	1.25	184
1984	1842	-62	39.64	9.03	48.67	0.81	62
1985	1780	-124	48.67	0.00	48.67	1.00	124
1986	1917	13	28.87	19.80	48.67	0.59	-13
1987	1890	-14	33.23	15.44	48.67	0.68	14
1988							
1989	1912	8	29.70	18.97	48.67	0.61	-8
1990							
1991							
1992							
1993	1782	-122	46.35	2.32	48.67	0.95	122
1994	1912	8	29.70	18.97	48.67	0.61	-8
1995	1805	-99	43.86	4.81	48.67	0.90	99
1996	1910	6	29.86	18.81	48.67	0.61	-6
1997	1715	-189	60.63	-11.96	48.67	1.25	189
1998	1935	31	25.53	23.14	48.67	0.52	-31
1999	2078	174	2.81	45.86	48.67	0.06	-174
2000	2052	148	5.83	42.84	48.67	0.12	-148
2001	1805	-99	43.87	4.80	48.67	0.90	99
2002	1960	56	20.67	28.00	48.67	0.42	-56
2003	1770	-134	47.72	0.95	48.67	0.98	134
2004	1782	-122	46.35	2.32	48.67	0.95	122
2005	1772	-132	47.72	0.95	48.67	0.98	132
2006	1910	6	29.86	18.81	48.67	0.61	-6
2007	1919	15	28.67	20.00	48.67	0.59	-15
2008	2015	111	11.10	37.57	48.67	0.23	-111
2009	1935	31	25.53	23.14	48.67	0.52	-31
2010	1902	-2	31.35	17.32	48.67	0.64	2
2011	2085	181	1.64	47.03	48.67	0.03	-181
MEAN	1888	-16	32.32	16.35		0.66	16



Photograph 28: McKenzie Glacier on 30 March 2011, resolution reduced to 200dpi.



Mountain glacier

GLACIER DATA		
Aspect	NW	
AREA	52.15	ha
Max Elev	2300	m
Min Elev	1920	m
Mean Elev	2110	m
Length	0.5	km
Elev Range	380	m
Gradient	0.76	

SNOWLINE DATA		
ELAo	2070	m
Max SL	2230	m, 1999
Min SL	1990	m, 1984
Mean SL	2065	m
SL Range	240	m
No. Surveys	27	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE	DEPARTURE		AREAS		ACCUM.	MASS
	ELEVATION	FROM ELAo	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
55	2100						
ELAo	2070	0	30.15	22.00	52.15	0.58	0
1977							
1978	2098	28	23.37	28.78	52.15	0.40	-28
1979							
1980	2050	-20	35.65	16.50	52.15	0.62	20
1981							
1982	2075	5	29.15	23.00	52.15	0.50	-5
1983	2032	-38	41.85	10.30	52.15	0.72	38
1984	1990	-80	49.05	3.10	52.15	0.85	80
1985	2014	-56	45.32	6.83	52.15	0.78	56
1986							
1987	2061	-9	32.71	19.44	52.15	0.56	9
1988							
1989	2075	5	29.15	23.00	52.15	0.50	-5
1990							
1991							
1992							
1993	1992	-78	48.97	3.18	52.15	0.85	78
1994	2045	-25	37.12	15.03	52.15	0.64	25
1995	2018	-52	44.31	7.84	52.15	0.77	52
1996	2072	2	29.97	22.18	52.15	0.52	-2
1997	2016	-54	44.85	7.30	52.15	0.77	54
1998	2075	5	27.05	18.45	45.50	0.47	-5
1999	2165	95	8.64	31.86	40.50	0.15	-95
2000	2115	45	15.49	23.01	38.50	0.27	-45
2001	2016	-54	32.40	6.10	38.50	0.56	54
2002	2103	33	22.77	15.33	38.10	0.39	-33
2003	2015	-55	36.60	1.50	38.10	0.63	55
2004	2016	-54	36.59	1.51	38.10	0.63	54
2005	2014	-56	36.61	1.49	38.10	0.63	56
2006	2085	15	22.60	15.50	38.10	0.39	-15
2007	2084	14	22.61	15.49	38.10	0.39	-14
2008	2145	75	9.10	28.50	37.60	0.24	-75
2009	2082	12	23.10	14.50	37.60	0.61	-12
2010	2072	2	26.30	11.30	37.60	0.70	-2
2011	2230	160	2.54	33.91	36.45	0.07	-160
MEAN	2065	-5	30.14	14.27		0.56	5





Photograph 29: Jackson Glacier on 30 March 2011, resolution reduced to 200dpi.

GLACIER DATA		
Aspect	W	
AREA	21.48	ha
Max Elev	2280	m
Min Elev	1800	m
Mean Elev	2040	m
Length	0.337	km
Elev Range	480	m
Gradient	1.42	

SNOWLINE DATA		
ELAo	1907	m
Max SL	2008	m, 1999
Min SL	1750	m, 1983
Mean SL	1898	m
SL Range	258	m
No. Surveys	30	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE	DEPARTURE	AREAS		ACCUM.	MASS
	ELEVATION	FROM ELAo	ACCUM.	ABL.	TOTAL	AREA RATIO
	m	m	ha	ha	ha	(AAR)
ELAo	1907	0	15.68	5.80	21.48	0.73
1977						
1978	1938	31	11.48	10.00	21.48	0.53
1979						
1980	1930	23	12.88	8.60	21.48	0.60
1981	1885	-22	17.35	4.13	21.48	0.81
1982	1951	44	9.00	12.48	21.48	0.42
1983	1750	-157	29.77	-8.29	21.48	1.39
1984	1828	-79	19.68	1.80	21.48	0.92
1985	1875	-32	17.88	3.60	21.48	0.83
1986	1935	28	11.98	9.50	21.48	0.56
1987	1905	-2	15.88	5.60	21.48	0.74
1988	1875	-32	17.88	3.60	21.48	0.83
1989	1898	-9	16.55	4.93	21.48	0.77
1990					no visit	
1991					no visit	
1992	1817	-90	19.88	1.60	21.48	0.93
1993	1765	-142	21.13	0.35	21.48	0.98
1994	1880	-27	17.68	3.80	21.48	0.82
1995	1755	-152	26.98	0.49	27.47	1.26
1996	1958	51	7.63	13.85	21.48	0.36
1997	1805	-102	20.19	1.29	21.48	0.94
1998	1940	33	11.15	10.33	21.48	0.52
1999	2008	101	1.96	19.52	21.48	0.09
2000	1992	85	2.86	18.62	21.48	0.13
2001	1798	-109	20.40	1.08	21.48	0.95
2002	1985	78	3.68	17.80	21.48	0.17
2003	1760	-147	21.48	0.00	21.48	1.00
2004					cloud	
2005	1795	-112	20.40	1.08	21.48	0.95
2006	1935	28	11.98	9.50	21.48	0.56
2007	1923	16	13.68	7.80	21.48	0.64
2008	1990	83	3.08	18.40	21.48	0.14
2009	1980	73	4.18	17.30	21.48	0.19
2010	1898	-9	16.55	4.93	21.48	0.77
2011	2200	293	0.00	21.48	21.48	0.00
MEAN	1898	-9	14.17	7.51		0.66
						9



Photograph 30: Jack Glacier on 30 March 2011, resolution reduced to 200dpi.

Rock glacier




GLACIER DATA		
Aspect	SE	
AREA	28.32	ha
Max Elev	2200	m
Min Elev	1760	m
Mean Elev	1980	m
Length	0.7	km
Elev Range	440	m
Gradient	0.629	

SNOWLINE DATA		
ELAo	1926	m
Max SL	2180	m, 2000
Min SL	1755	m, 1984
Mean SL	1913	m
SL Range	425	m
No. Surveys	24	

MEASUREMENTS



Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1926	0	6.52	21.80	28.32	0.23	0
1977							
1978							
1979							
1980							
1981							
1982							
1983							
1984	1755	-171	20.32	8.00	28.32	0.72	171
1985							
1986	1835	-91	12.33	15.99	28.32	0.44	91
1987							
1988	1907	-19	7.92	20.40	28.32	0.28	19
1989	1972	46	4.41	23.91	28.32	0.16	-46
1990							
1991							
1992	1850	-76	11.32	17.00	28.32	0.40	76
1993	1785	-141	16.69	11.63	28.32	0.59	141
1994	1850	-76	11.32	17.00	28.32	0.40	76
1995	1770	-156	18.39	9.93	28.32	0.65	156
1996	1855	-71	11.08	17.24	28.32	0.39	71
1997	1842	-84	11.75	16.57	28.32	0.41	84
1998	1889	-37	8.93	19.39	28.32	0.32	37
1999	2125	199	0.42	27.90	28.32	0.01	-199
2000	2130	204	0.37	27.95	28.32	0.01	-204
2001	1780	-146	17.22	11.10	28.32	0.61	146
2002	2115	189	0.52	27.80	28.32	0.02	-189
2003	1795	-131	15.72	12.60	28.32	0.56	131
2004	1855	-71	11.08	17.24	28.32	0.39	71
2005	1800	-126	15.72	12.60	28.32	0.56	126
2006	1895	-31	8.52	19.80	28.32	0.30	31
2007	1910	-16	7.62	20.70	28.32	0.27	16
2008	2120	194	0.42	27.90	28.32	0.01	-194
2009	2000	74	3.22	25.10	28.32	0.11	-74
2010	1885	-41	9.12	19.20	28.32	0.32	41
2011	2180	254	0.00	28.32	28.32	0.00	-254
MEAN	1913	-14	9.76	18.56		0.34	14



Photograph 31: Mt St Mary on 30 March 2011, resolution reduced to 200dpi.

No. 711B/12			THURNEYSTON GL.			NZMS 260 sheet G38	
			Mountain gl, reliability C				
	GLACIER DATA				SNOWLINE DATA		
	Aspect	S					
	AREA	117.8	ha		ELA	1970	m
	Max Elev	2450	m		Max SL	2150	m, 2000
	Min Elev	1720	m		Min SL	1865	m, 1983
	Mean Elev	2085	m		Mean SL	1962	m
	Length	1.23	km		SL Range	285	m
	Elev Range	730	m		No. Surveys	30	
	Gradient	0.59					
MEASUREMENTS		Digitised values shaded					
YEAR	SNOWLINE	DEPARTURE		AREAS		ACCUM.	MASS
	ELEVATION	FROM ELA	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	1970	0	84.60	33.20	117.80	0.72	0
1977	1930	-40	95.30	22.50	117.80	0.81	40
1978							
1979							
1980	1926	-44	98.00	19.80	117.80	0.83	44
1981	1943	-27	92.80	25.00	117.80	0.79	27
1982							
1983	1865	-105	109.80	8.00	117.80	0.93	105
1984	1882	-88	107.00	10.80	117.80	0.91	88
1985	1905	-65	103.40	14.40	117.80	0.88	65
1986	1938	-32	94.08	23.72	117.80	0.80	32
1987	1918	-52	100.30	17.50	117.80	0.85	52
1988	1950	-20	90.80	27.00	117.80	0.77	20
1989	1970	0	84.60	33.20	117.80	0.72	0
1990							
1991							
1992	1930	-40	95.30	22.50	117.80	0.81	40
1993	1910	-60	101.90	15.90	117.80	0.87	60
1994	1904	-66	103.40	14.40	117.80	0.88	66
1995	1868	-102	109.40	8.40	117.80	0.93	102
1996	1938	-32	94.08	23.72	117.80	0.80	32
1997	1900	-70	103.85	13.95	117.80	0.88	70
1998	1965	-5	85.28	32.52	117.80	0.72	5
1999	2112	142	38.69	79.11	117.80	0.33	-142
2000	2132	162	31.67	86.13	117.80	0.27	-162
2001	1878	-92	107.84	9.96	117.80	0.92	92
2002	2105	135	40.80	77.00	117.80	0.35	-135
2003	1873	-97	108.60	9.20	117.80	0.92	97
2004	1935	-35	95.30	22.50	117.80	0.81	35
2005	1900	-70	103.85	13.95	117.80	0.88	70
2006	1980	10	81.80	36.00	117.80	0.69	-10
2007	1963	-7	85.70	32.10	117.80	0.73	7
2008	2110	140	39.80	78.00	117.80	0.34	-140
2009	2100	130	42.80	75.00	117.80	0.36	-130
2010	1975	5	82.80	35.00	117.80	0.70	-5
2011	2150	180	24.80	93.00	117.80	0.21	-180
MEAN	1962	-8	85	32.68		0.72	8



Photograph 32: Thurneyson Glacier on 30 March 2011, resolution reduced to 200dpi.

Mountain glacier



GLACIER DATA		
Aspect	S	
AREA	249.5	ha
Max Elev	2390	m
Min Elev	1655	m
Mean Elev	2023	m
Length	2.69	km
Elev Range	735	m
Gradient	0.27	

SNOWLINE DATA		
ELAo	1935	m
Max SL	2285	m, 1999
Min SL	1750	m, 1993
Mean SL	1909	m
SL Range	535	m
No. Surveys	30	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1935	0	125.6	128.0	253.6	0.50	0
1977							
1978	1960	25	98.7	154.9	253.6	0.39	-25
1979							
1980	1846	-89	193.5	60.1	253.6	0.76	89
1981	1855	-80	188.5	65.1	253.6	0.74	80
1982	1971	36	80.2	173.4	253.6	0.32	-36
1983	1794	-141	221.3	32.3	253.6	0.87	141
1984	1800	-135	218.4	35.2	253.6	0.86	135
1985	1796	-139	220.5	33.1	253.6	0.87	139
1986	1842	-93	195.5	58.1	253.6	0.77	93
1987	1828	-107	205.6	48.0	253.6	0.81	107
1988							
1989	1918	-17	140.4	113.2	253.6	0.55	17
1990							
1991							
1992	1851	-84	190.5	63.1	253.6	0.75	84
1993	1750	-185	234.7	18.9	253.6	0.93	185
1994	1790	-145	222.5	31.1	253.6	0.88	145
1995	1777	-158	227.6	26.0	253.6	0.90	158
1996	1962	27	89.9	163.7	253.6	0.35	-27
1997	1779	-156	226.3	27.0	253.3	0.89	156
1998	1982	47	67.0	186.1	253.1	0.26	-47
1999	2280	345	8.2	244.6	252.8	0.03	-345
2000	2155	220	29.2	223.4	252.5	0.12	-220
2001	1770	-165	228.7	20.8	252.2	0.91	165
2002	2050	115	203.0	49.0	252.0	0.81	-115
2003	1794	-141	222.7	29.0	251.7	0.88	141
2004	1780	-155	227.4	24.0	251.4	0.90	155
2005	1780	-155	227.4	23.7	251.1	0.91	155
2006	1870	-65	179.1	71.8	250.9	0.71	65
2007	1815	-120	211.8	38.8	250.6	0.85	120
2008	2270	335	8.4	241.1	249.5	0.03	-335
2009	1990	55	62.8	186.7	249.5	0.25	-55
2010	1930	-5	130.1	119.4	249.5	0.52	5
2011	2285	350	3.5	246.0	249.5	0.01	-350
MEAN	1909	-26	159	88		0.63	26



Photograph 33: Brewster Glacier on 30 March 2011, resolution reduced to 200dpi.

Moutain glacier



GLACIER DATA		
Aspect	SE	
AREA	31.55	ha
Max Elev	1860	m
Min Elev	1570	m
Mean Elev	1715	m
Length	0.54	km
Elev Range	290	m
Gradient	0.54	

SNOWLINE DATA		
ELAo	1673	m
Max SL	1858	m, 2000
Min SL	1515	m, 1995
Mean SL	1664	m
SL Range	343	m
No. Surveys	29	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE	DEPARTURE		AREAS		ACCUM.	MASS
	ELEVATION	FROM ELAo	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	1673	0	21.05	5.00	26.05	0.81	0
1977	1587	-86	30.86	0.69	31.55	0.98	86
1978	1730	57	18.53	13.02	31.55	0.59	-57
1979							
1980	1650	-23	27.25	4.30	31.55	0.86	23
1981	1606	-67	29.85	1.70	31.55	0.95	67
1982	1676	3	25.33	6.22	31.55	0.80	-3
1983	1538	-135	31.55	0.00	31.55	1.00	135
1984							
1985	1620	-53	29.10	2.45	31.55	0.92	53
1986	1660	-13	26.55	5.00	31.55	0.84	13
1987	1678	5	25.07	6.48	31.55	0.79	-5
1988							
1989	1663	-10	26.32	5.23	31.55	0.83	10
1990							
1991							
1992							
1993	1535	-138	31.55	0.00	31.55	1.00	138
1994	1640	-33	27.95	3.60	31.55	0.89	33
1995	1515	-158	31.55	0.00	31.55	1.00	158
1996	1712	39	21.20	10.35	31.55	0.67	-39
1997	1567	-106	31.55	0.00	31.55	1.00	106
1998	1656	-17	26.87	4.68	31.55	0.85	17
1999	1805	132	5.46	26.09	31.55	0.17	-132
2000	1850	177	0.95	25.10	26.05	0.04	-177
2001	1590	-83	25.09	0.96	26.05	0.96	83
2002	1815	142	4.45	21.60	26.05	0.17	-142
2003	1570	-103	26.05	0.00	26.05	1.00	103
2004	1565	-108	26.05	0.00	26.05	1.00	108
2005	1565	-108	26.05	0.00	26.05	1.00	108
2006	1725	52	16.85	9.20	26.05	0.65	-52
2007	1607	-66	24.65	1.40	26.05	0.95	66.00
2008	1800	127	6.35	19.70	26.05	0.24	-127.00
2009	1790	117	7.65	18.40	26.05	0.29	-117.00
2010	1677	4	20.45	5.60	26.05	0.79	-4.00
2011	1858	185	0.29	25.76	26.05	0.01	-185.00
MEAN	1664	-9	21.77	7.50		0.73	9



Photograph 34: Stuart Glacier on 30 March 2011, resolution reduced to 200dpi.




GLACIER DATA		
Aspect	S	
AREA	52.53	ha
Max Elev	1880	m
Min Elev	1610	m
Mean Elev	1745	m
Length	0.57	km
Elev Range	270	m
Gradient	0.47	

SNOWLINE DATA		
ELAo	1730	m
Max SL	1878	m, 1999
Min SL	1550	m, 1995
Mean SL	1716	m
SL Range	328	m
No. Surveys	29	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE ELEVATION m	DEPARTURE FROM ELAo m	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1730	0	39.23	13.30	52.53	0.75	0
1977							
1978	1738	8	37.09	15.44	52.53	0.71	-8
1979							
1980	1652	-78	50.51	2.02	52.53	0.96	78
1981	1681	-49	48.60	3.93	52.53	0.93	49
1982	1781	51	22.41	30.12	52.53	0.43	-51
1983	1560	-170	65.66	0.00	65.66	1.25	170
1984							
1985	1666	-64	49.63	2.90	52.53	0.94	64
1986	1768	38	27.53	25.00	52.53	0.52	-38
1987	1615	-115	52.33	0.20	52.53	1.00	115
1988	1772	42	24.63	27.90	52.53	0.47	-42
1989	1764	34	28.49	24.04	52.53	0.54	-34
1990							
1991							
1992							
1993	1555	-175	69.94	0.00	69.94	1.33	175
1994	1610	-120	52.53	0.00	52.53	1.00	120
1995	1550	-180	73.50	0.00	73.50	1.40	180
1996	1775	45	24.47	28.06	52.53	0.47	-45
1997	1645	-85	69.00	1.65	70.65	1.31	85
1998	1800	70	14.45	38.08	52.53	0.28	-70
1999	1875	145	0.94	51.59	52.53	0.02	-145
2000	1870	140	1.47	51.06	52.53	0.03	-140
2001	1640	-90	51.09	1.44	52.53	0.97	90
2002	1872	142	1.05	51.48	52.53	0.02	-142
2003	1607	-123	52.53	0.00	52.53	1.00	123
2004	1608	-122	52.53	0.00	52.53	1.00	122
2005	1612	-118	51.53	1.00	52.53	0.98	118
2006	1780	50	22.40	30.13	52.53	0.43	-50
2007	1700	-30	46.43	6.10	52.53	0.88	30
2008	1860	130	2.53	50.00	52.53	0.05	-130
2009	1810	80	11.93	40.60	52.53	0.23	-80
2010	1710	-20	44.53	8.00	52.53	0.85	20
2011	1878	148	0.53	52.00	52.53	0.01	-148
MEAN	1716	-14	36.22	18.72		0.69	20



Photograph 35: Lindsay Glacier on 30 March 2011, resolution reduced to 200dpi.

Cirque glacier


GLACIER DATA		
Aspect	SE	
AREA	18.22	ha
Max Elev	2150	m
Min Elev	1840	m
Mean Elev	1995	m
Length	0.4	km
Elev Range	310	m
Gradient	0.775	

SNOWLINE DATA		
ELAo	1987	m
Max SL	2122	m, 1999
Min SL	1888	m, 1995
Mean SL	1984	m
SL Range	234	m
No. Surveys	26	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	ACCUM. TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1987	0	12.02	6.20	18.22	0.66	0
1977							
1978							
1979							
1980	1916	-71	34.40	0.00	34.40	>1	71
1981							
1982	2022	35	14.75	3.47	18.22	0.81	-35
1983							
1984	1891	-96	42.80	0.00	42.80	>1	96
1985							
1986	1930	-57	30.40	0.00	30.40	>1	57
1987	1902	-85	39.00	0.00	39.00	>1	85
1988	2022	35	14.75	3.47	18.22	0.81	-35
1989	2032	45	6.62	11.60	18.22	0.36	-45
1990							
1991							
1992							
1993	1894	-93	37.80	0.00	37.80	>1	93
1994	1900	-87	17.58	0.64	18.22	0.96	87
1995	1888	-99	44.36	0.00	44.36	>1	99
1996	1894	-93	41.70	0.00	41.70	>1	93
1997	1890	-97	43.25	0.00	43.25	>1	97
1998	2098	111	1.77	16.45	18.22	0.10	-111
1999	2122	135	0.74	17.48	18.22	0.04	-135
2000	2108	121	1.28	16.94	18.22	0.07	-121
2001	1895	-92	17.42	0.80	18.22	0.96	92
2002	2112	125	1.13	17.09	18.22	0.06	-125
2003	1962	-25	15.40	2.82	18.22	0.85	25
2004	1892	-95	18.22	0.00	18.22	1.00	95
2005	1890	-97	18.22	0.00	18.22	1.00	97
2006	2032	45	6.62	11.60	18.22	0.36	-45
2007	2000	13	10.22	8.00	18.22	0.56	-13
2008	2105	118	1.42	16.80	18.22	0.08	-118
2009	2032	45	6.62	11.60	18.22	0.36	-45
2010	2032	45	6.62	11.60	18.22	0.36	-45
2011	2122	135	0.74	17.48	18.22	0.04	-135
MEAN	1984	-3	18.22	6.46		0.49	3



Photograph 36: Fog Peak on 30 March 2011, resolution reduced to 200dpi.



GLACIER DATA		
Aspect	W	
AREA	53.13	ha
Max Elev	2210	m
Min Elev	2000	m
Mean Elev	2105	m
Length	0.73	km
Elev Range	210	m
Gradient	0.29	

SNOWLINE DATA		
ELAo	2092	m
Max SL	2240	m, 1999
Min SL	2004	m, 1997
Mean SL	2081	m
SL Range	236	m
No. Surveys	30	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE	DEPARTURE		AREAS		ACCUM.	MASS
	ELEVATION	FROM ELAo	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	2092	0	31.88	21.25	53.13	0.60	0
1977							
1978	2156	64	6.86	46.27	53.13	0.13	-64
1979							
1980	2024	-68	48.36	4.77	53.13	0.91	68
1981	2158	66	6.42	46.71	53.13	0.12	-66
1982	2038	-54	45.43	7.70	53.13	0.86	54
1983	2033	-59	53.13	0.00	53.13	1.00	59
1984	2024	-68	48.33	4.80	53.13	0.91	68
1985	2020	-72	49.13	4.00	53.13	0.92	72
1986	2036	-56	45.93	7.20	53.13	0.86	56
1987	2025	-67	48.19	4.94	53.13	0.91	67
1988	2037	-55	45.93	7.20	53.13	0.86	55
1989	2103	11	28.18	24.95	53.13	0.53	-11
1990							
1991							
1992							
1993	2058	-34	40.69	12.44	53.13	0.77	34
1994	2034	-58	46.13	7.00	53.13	0.87	58
1995	2030	-62	47.13	6.00	53.13	0.89	62
1996	2057	-35	41.04	12.09	53.13	0.77	35
1997	2020	-72	64.34	-11.21	53.13	1.21	72
1998	2158	66	6.42	46.71	53.13	0.12	-66
1999	2240	148	53.13	0.00	53.13	1.00	-148
2000	2160	68	6.13	47.00	53.13	0.12	-68
2001	2004	-88	52.73	0.40	53.13	0.99	88
2002	2120	28	21.83	31.30	53.13	0.41	-28
2003	2041	-51	44.73	8.40	53.13	0.84	51
2004	2034	-58	53.13	0.00	53.13	1.00	58
2005	2020	-72	50.13	3.00	53.13	0.94	72
2006	2160	68	6.13	47	53.13	0.12	-68
2007	2090	-2	32.13	21.00	53.13	0.60	2
2008	2158	66	6.73	46.40	53.13	0.13	-66
2009	2156	64	6.86	46.27	53.13	0.13	-64
2010	2156	64	6.86	46.27	53.13	0.13	-64
2011	>2210		0.00	53.13	53.13	0.00	
MEAN	2081	-11	35.90	17.23		0.68	11



Photograph 37: Snowy Peak on 30 March 2011, resolution reduced to 200dpi.

Small cirque




GLACIER DATA		
Aspect	SE	
AREA	17.13	ha
Max Elev	1600	m
Min Elev	1400	m
Mean Elev	1500	m
Length	0.3	km
Elev Range	200	m
Gradient	0.67	

SNOWLINE DATA		
ELAo	1472	m
Max SL	1660	m, 2000
Min SL	1366	m, 1995
Mean SL	1453	m
SL Range	294	m
No. Surveys	28	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	ACCUM. TOTAL ha	MASS AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1472	0	2.83	14.30	17.13	0.17	0
1977							
1978	1442	-30	7.05	10.08	17.13	0.41	30
1979							
1980							
1981	1413	-59	13.66	3.47	17.13	0.80	59
1982	1424	-48	10.74	6.39	17.13	0.63	48
1983	1372	-100	42.80	0.00	42.8	2.50	100
1984							
1985	1423	-49	10.84	6.29	17.13	0.63	49
1986	1525	53	9.80	7.33	17.13	0.57	-53
1987	1422	-50	11.06	6.07	17.13	0.65	50
1988	1500	28	6.43	10.70	17.13	0.38	-28
1989	1429	-43	9.80	7.33	17.13	0.57	43
1990							
1991							
1992							
1993	1375	-97	35.98	0.00	35.98	2.10	97
1994	1400	-72	17.13	0.00	17.13	1.00	72
1995	1366	-106	54.48	0.00	54.48	3.18	106
1996	1390	-82	22.00	0.00	22	1.28	82
1997	1395	-77	19.50	0.00	19.5	1.14	77
1998	1417	-55	12.46	4.67	17.13	0.73	55
1999	1650	178	0.00	17.13	17.13	0.00	-178
2000	1660	188	0.00	8.74	8.74	0.00	-188
2001	1420	-52	9.15	3.50	12.65	0.72	52
2002	1625	153	0.00	8.74	8.74	0.00	-153
2003	1422	-50	8.14	0.60	8.74	0.93	50
2004	1380	-92	8.74	0.00	8.74	1.00	92
2005	1374	-98	35.98	0.00	35.98	1.00	98
2006	1515	43	0.74	8.00	8.74	0.08	-43
2007	1455	-17	3.74	5.00	8.74	0.43	17
2008	1520	48	0.44	8.30	8.74	0.05	-48
2009	1520	48	0.44	8.30	8.74	0.05	-48
2010	1398	-74	8.74	0.00	8.74	1.00	74
2011	>1660		0.00	8.74	8.74	0.00	
MEAN	1453	-19	12.85	4.98		0.78	17



Photograph 38: Mt Caria on 30 March 2011, resolution reduced to 200dpi.



Cirque glacier

GLACIER DATA		
Aspect	SW	
AREA	67.15	ha
Max Elev	1900	m
Min Elev	1550	m
Mean Elev	1725	m
Length	0.875	km
Elev Range	350	m
Gradient	0.4	

SNOWLINE DATA		
ELAo	1693	m
Max SL	1890	m, 1999
Min SL	1561	m, 1995
Mean SL	1682	m
SL Range	329	m
No. Surve	27	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE	DEPARTURE		AREAS		ACCUM.	MASS
	ELEVATION	FROM ELAo	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	1693	0	35.95	31.2	67.15	0.54	0
1977							
1978							
1979							
1980							
1981	1604	-89	57.53	9.62	67.15	0.86	89
1982	1735	42	25.85	41.30	67.15	0.38	-42
1983	1582	-111	61.52	5.63	67.15	0.92	111
1984							
1985	1629	-64	48.89	18.26	67.15	0.73	64
1986	1725	32	27.95	39.20	67.15	0.42	-32
1987	1622	-71	51.05	16.10	67.15	0.76	71
1988	1685	-8	35.15	32.00	67.15	0.52	8
1989	1642	-51	44.15	23.00	67.15	0.66	51
1990							
1991							
1992							
1993	1575	-118	62.90	4.25	67.15	0.94	118
1994	1634	-59	46.87	20.28	67.15	0.70	59
1995	1561	-132	65.17	1.98	67.15	0.97	132
1996	1632	-61	47.54	19.61	67.15	0.71	61
1997	1580	-113	62.00	5.15	67.15	0.92	113
1998	1780	87	22.91	35.49	58.4	0.39	-87
1999	1890	197	2.11	51.44	53.55	0.04	-197
2000	1845	152	7.28	44.32	51.6	0.14	-152
2001	1620	-73	45.36	8.80	54.16	0.84	73
2002	1825	132	10.79	46.63	51.6	0.21	-132
2003	1612	-81	42.40	11.80	54.2	0.78	81
2004	1578	-115	49.80	4.40	54.2	0.92	115
2005	1580	-113	49.80	4.40	54.2	0.92	113
2006	1690	-3	31.95	22.21	54.16	0.59	3
2007	1661	-32	35.06	19.10	54.16	0.65	32
2008	1832	139	8.50	45.66	54.16	0.16	-139
2009	1805	112	14.96	39.20	54.16	0.28	-112
2010	1632	-61	41.66	12.50	54.16	0.77	61
2011	1870	177	4.36	49.80	54.16	0.08	-177
Mean	1682	-11	39.27	22.11		0.63	11





Photograph 39: Findlay Glacier on 30 March 2011, resolution reduced to 200dpi.

GLACIER DATA		
Aspect	S	
AREA	191.2	ha
Max Elev	2200	m
Min Elev	1500	m
Mean Elev	1850	m
Length	2.63	km
Elev Range	700	m
Gradient	0.267	

SNOWLINE DATA		
ELAo	1824	m
Max SL	2005	m, 1999
Min SL	1635	m, 1995
Mean SL	1819	m
SL Range	370	m
No. Surveys	28	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1824	0	127.90	85.70	213.60	0.60	0
1977							
1978	1903	79	91.29	122.31	213.60	0.43	-79
1979							
1980	1808	-16	131.60	82.00	213.60	0.62	16
1981	1778	-46	139.20	74.40	213.60	0.65	46
1982	1858	34	112.40	101.20	213.60	0.53	-34
1983	1762	-62	142.67	70.93	213.60	0.67	62
1984	1765	-59	141.60	72.00	213.60	0.66	59
1985	1702	-122	156.60	57.00	213.60	0.73	122
1986	1863	39	110.10	103.50	213.60	0.52	-39
1987	1843	19	117.60	96.00	213.60	0.55	-19
1988							
1989	1794	-30	134.60	79.00	213.60	0.63	30
1990							
1991							
1992							
1993							
1994	1783	-41	151.40	55.00	206.40	0.73	41
1995	1635	-189	187.60	18.30	205.90	0.91	189
1996	1808	-16	147.17	58.23	205.40	0.72	16
1997	1745	-79	160.20	44.10	204.30	0.78	79
1998	1880	56	105.80	95.60	201.40	0.53	-56
1999	1955	131	62.07	136.03	198.10	0.31	-131
2000	1935	111	71.34	124.66	196.00	0.36	-111
2001	1748	-76	149.44	45.36	194.80	0.77	76
2002	1910	86	82.82	111.18	194.00	0.43	-86
2003	1661	-163	166.80	26.50	193.30	0.86	163
2004	1665	-159	162.60	30.00	192.60	0.84	159
2005	1670	-154	161.00	31.00	192.00	0.84	154
2006	1900	76	86.20	105.00	191.20	0.45	-76
2007	1850	26	109.50	80.40	189.90	0.58	-26
2008	1910	86	81.20	107.30	188.50	0.43	-86
2009	1910	86	79.80	107.30	187.10	0.43	-86
2010	1880	56	95.20	91.50	186.70	0.51	-56
2011	2005	181	35.88	148.52	184.40	0.19	-181
Mean	1819	-5	123.62	78.73		0.59	5



Photograph 40: Park Pass Glacier on 30 March 2011, resolution reduced to 200dpi.

Glacierette




GLACIER DATA		
Aspect	SE	
AREA	27.47	ha
Max Elev	2220	m
Min Elev	1680	m
Mean Elev	1950	m
Length	0.5	km
Elev Range	540	m
Gradient	1.08	

SNOWLINE DATA		
ELAo	1945	m
Max. SL	2215	m, 1998
Min. SL	1630	m, 1995
Mean SL	1934	m
SL Range	585	m
No. surveys	24	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	ACCUM. TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1945	0	17.67	9.80	27.47	0.64	0
1977							
1978							
1979							
1980							
1981							
1982							
1983							
1984	1680	-265	33.82	0.00	33.82	1.23	265
1985							
1986	1892	-53	33.82	0.00	33.82	1.23	53
1987							
1988	2050	105	9.47	18.00	27.47	0.34	-105
1989	2036	91	10.84	16.63	27.47	0.39	-91
1990							
1991							
1992							
1993	1670	-275	39.50	0.00	39.50	1.44	275
1994	1850	-95	23.40	4.07	27.47	0.85	95
1995	1630	-315	43.61	0.00	43.61	1.59	315
1996	1782	-163	27.47	0.00	27.47	1.00	163
1997	1633	-312	36.80	0.00	36.80	1.34	312
1998	2050	105	0.92	26.55	27.47	0.03	-105
1999	2215	270	0.95	26.52	27.47	0.03	-270
2000	2200	255	1.46	14.75	16.21	0.09	-255
2001	1665	-280	27.47	0.00	27.47	1.00	280
2002	2095	150	2.17	14.04	16.21	0.13	-150
2003	1740	-205	15.51	0.70	16.21	0.96	205
2004	1690	-255	27.47	0.00	27.47	1.00	255
2005	1660	-285	27.47	0.00	27.47	1.00	285
2006	2070	125	1.46	14.75	16.21	0.09	-125
2007	2090	145	2.20	14.01	16.21	0.14	-145
2008	2190	245	1.50	14.62	16.12	0.09	-245
2009	2189	244	1.60	14.52	16.12	0.10	-244
2010	2180	235	2.20	13.92	16.12	0.14	-235
2011	2215	270	0.90	15.22	16.12	0.06	-270
MEAN	1934	-11	16.17	9.06		0.62	11



Photograph 43: Mt Larkins on 30 March 2011, resolution reduced to 200dpi.

Mountain glacier



GLACIER DATA		
Aspect	SE	
AREA	29.7	ha
Max Elev	2180	m
Min Elev	1660	m
Mean Elev	1920	m
Length	0.94	km
Elev Range	520	m
Gradient	0.55	

SNOWLINE DATA		
ELAo	1783	m
Max. SL	2050	m, 1999
Min. SL	1610	m, 1995
Mean SL	1762	m
SL Range	440	m
No. surveys	29	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1783	0	12.20	17.50	29.7	0.41	0
1977	1740	-43	20.90	8.80	29.7	0.70	43
1978	1884	101	4.90	24.80	29.7	0.16	-101
1979							
1980							
1981	1763	-20	15.70	14.00	29.7	0.53	20
1982	1780	-3	12.70	17.00	29.7	0.43	3
1983	1620	-163	34.50	0.00	29.7	1.00	163
1984	1620	-163	31.80	0.00	29.7	1.00	163
1985	1610	-173	37.30	0.00	29.7	1.00	173
1986	1770	-13	14.08	15.62	29.7	0.47	13
1987	1763	-20	15.70	14.00	29.7	0.53	20
1988							
1989	1753	-30	17.40	12.30	29.7	0.59	30
1990							
1991							
1992							
1993	1675	-108	17.20	12.50	29.7	0.58	108
1994	1728	-55	23.40	6.30	29.7	0.79	55
1995	1630	-153	40.17	0.00	29.7	1.00	153
1996	1680	-103	29.10	0.60	29.7	0.98	103
1997	1670	-113	29.01	0.69	29.7	0.98	113
1998	1778	-5	13.16	16.54	29.7	0.44	5
1999	2010	227	0.97	28.73	29.7	0.03	-227
2000	1965	182	1.85	27.85	29.7	0.06	-182
2001	1665	-118	29.20	0.50	29.7	0.98	118
2002	1870	87	5.50	24.20	29.7	0.19	-87
2003	1666	-117	28.80	0.90	29.7	0.97	117
2004	1675	-108	29.06	0.64	29.7	0.98	108
2005	1665	-118	28.80	0.90	29.7	0.97	118
2006	1800	17	10.10	19.60	29.7	0.34	-17
2007	1775	-8	13.50	16.20	29.7	0.45	8
2008	1950	167	2.30	27.40	29.7	0.08	-167
2009	1775	-8	13.50	16.20	29.7	0.45	8
2010	1755	-28	17.70	12.00	29.7	0.60	28
2011	2050	267	0.60	29.10	29.7	0.02	-267
MEAN	1762	-21	18.58	11.98		0.60	21



Photograph 45: Bryant Glacier on 30 March 2011, resolution reduced to 200dpi.



Cirque glacier

GL DATA		
Aspect	S	
AREA	22.91	ha
Max Elev	1830	m
Min Elev	1530	m
Mean Elev	1680	m
Length	0.75	km
Elev Range	300	m
Gradient	0.4	

SNOWLINE DATA		
ELAo	1648	m
Max SL	1830	m, 1999
Min SL	1555	m, 1995
Mean SL	1632	m
SL Range	275	m
No. surveys	26	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE	DEPARTURE	AREAS		ACCUM.	MASS	
	ELEVATION	FROM ELAo	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	1648	0	7.72	10.50	18.22	0.42	0
1977							
1978	1643	-5	9.32	13.59	22.91	0.41	5
1979							
1980							
1981							
1982							
1983	1560	-88	17.65	3.70	21.35	0.77	88
1984	1595	-53	13.20	8.00	21.2	0.58	53
1985	1595	-53	13.10	8.00	21.1	0.57	53
1986	1649	1	8.57	12.23	20.8	0.37	-1
1987	1625	-23	11.48	8.72	20.2	0.50	23
1988							
1989	1612	-36	9.97	10.13	20.1	0.44	36
1990							
1991							
1992							
1993	1564	-84	15.43	4.10	19.53	0.67	84
1994	1596	-52	11.06	8.20	19.26	0.48	52
1995	1555	-93	15.87	3.12	18.99	0.69	93
1996	1584	-64	12.06	6.64	18.7	0.53	64
1997	1593	-55	12.16	6.28	18.44	0.53	55
1998	1621	-27	11.91	4.99	16.9	0.52	27
1999	1830	182	0.00	22.91	15.3	0.00	-182
2000	1785	137	0.77	22.14	14.79	0.05	-137
2001	1593	-55	9.51	5.28	14.79	0.64	55
2002	1685	37	4.35	13.87	14.5	0.30	-37
2003	1583	-65	12.20	2.00	14.2	0.86	65
2004	1590	-58	11.74	2.31	14.05	0.84	58
2005	1580	-68	11.99	1.96	13.95	0.86	68
2006	1645	-3	6.90	7.00	13.9	0.50	3
2007	1613	-35	9.80	4.10	13.9	0.71	35
2008	1670	22	3.70	10.20	13.9	0.27	-22
2009	1675	27	3.30	10.60	13.9	0.24	-27
2010	1615	-33	9.90	4.00	13.9	0.71	33
2011	1780	132	0.89	12.81	13.7	0.06	-132
MEAN	1632	-16	9.49	8.34		0.50	16



Photograph 47: Ailsa Glacier on 12 March 2011, resolution reduced to 200dpi.



GLACIER DATA		
Aspect	SE	
AREA	53.72	ha
Max Elev	1860	m
Min Elev	1470	m
Mean Elev	1665	m
Length	0.75	km
Elev Range	390	m
Gradient	0.52	

SNOWLINE DATA		
ELAo	1593	m
Max SL	1820	m, 2000
Min SL	1471	m, 1995
Mean SL	1582	m
SL Range	349	m
No. surveys	28	

MEASUREMENTS



Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	AREAS ACCUM.	AREAS ABL.	ACCUM. TOTAL	MASS AREA RATIO	MASS BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	1593	0	14.67	21.00	35.67	0.41	0
1977	1615	22	14.72	39.00	53.72	0.27	-22
1978	1638	45	12.11	41.61	53.72	0.23	-45
1979							
1980	1529	-64	33.40	20.32	53.72	0.62	64
1981	1531	-62	32.44	21.28	53.72	0.60	62
1982	1610	17	15.50	38.22	53.72	0.29	-17
1983	1478	-115	51.31	2.41	53.72	0.96	115
1984							
1985	1540	-53	30.24	23.48	53.72	0.56	53
1986							
1987	1555	-38	26.32	27.40	53.72	0.49	38
1988							
1989	1534	-59	31.84	20.60	52.44	0.59	59
1990							
1991							
1992							
1993	1485	-108	48.12	2.68	50.80	0.90	108
1994	1529	-64	33.40	16.58	49.98	0.62	64
1995	1471	-122	54.00	0.00	49.00	1.01	122
1996	1507	-86	40.12	7.58	47.70	0.75	86
1997	1520	-73	36.19	10.01	46.20	0.67	73
1998	1559	-34	25.42	19.29	44.71	0.47	34
1999	1802	209	0.77	41.73	42.50	0.01	-209
2000	1810	217	0.49	40.49	40.98	0.01	-217
2001	1515	-78	33.93	6.80	40.73	0.83	78
2002	1635	42	8.83	31.65	40.48	0.22	-42
2003	1525	-68	32.63	7.60	40.23	0.81	68
2004	1485	-108	37.88	2.10	39.98	0.95	108
2005	1485	-108	37.63	2.10	39.73	0.95	108
2006	1610	17	22.18	17.30	39.48	0.56	-17
2007	1550	-43	28.60	9.50	38.10	0.75	43
2008	1645	52	19.00	18.00	37.00	0.51	-52
2009	1790	197	7.10	28.55	35.65	0.20	-197
2010	1515	-78	33.30	0.70	34.00	0.98	78
2011	1820	227	0.40	33.60	34.00	0.01	-227
MEAN	1582	-11	26.71	18.95		0.57	11



Photograph 48: Mt Gunn on 12 March 2011, resolution reduced to 200dpi.

No. 797G/33			Mt. GENDARME			NZMS sheet D40 & 41	
			Mountain gl, reliability B				
	GLACIER DATA				SNOWLINE DATA		
	Aspect	S					
	Area	49.35	ha		ELA	1616	m
	Max Elev	1900	m		Max SL	1804	m, 1999
	Min Elev	1440	m		Min SL	1418	m, 1995
	Mean Elev	1670	m		Mean SL	1587	m
	Length	0.525	km		SL range	386	m
	Elev Range	460	m		No. surveys	26	
	Gradient	0.88					
MEASUREMENTS		Digitised values shaded					
YEAR	SNOWLINE	DEPARTURE		AREAS		ACCUM.	MASS
	ELEVATION	FROM ELA	ACCUM.	ABL.	TOTAL	AREA RATIO	BALANCE
	m	m	ha	ha	ha	(AAR)	INDEX
ELAo	1616	0	31.85	17.50	49.35	0.65	0
1977							
1978							
1979							
1980							
1981	1570	-46	38.36	10.99	49.35	0.78	46
1982	1573	-43	37.85	11.50	49.35	0.77	43
1983	1480	-136	55.97	0.00	55.97	1.13	136
1984							
1985	1522	-94	44.53	4.82	49.35	0.90	94
1986	1675	59	21.95	27.40	49.35	0.44	-59
1987	1650	34	25.87	23.48	49.35	0.52	-34
1988							
1989	1580	-36	35.50	13.85	49.35	0.72	36
1990							
1991							
1992							
1993	1502	-114	46.42	2.93	49.35	0.94	114
1994	1552	-64	40.55	8.80	49.35	0.82	64
1995	1418	-198	58.67	0.00	58.67	1.19	198
1996	1516	-100	45.20	4.15	49.35	0.92	100
1997	1490	-126	47.25	2.10	49.35	0.96	126
1998	1648	32	26.45	22.90	49.35	0.54	-32
1999	1804	188	2.33	47.02	49.35	0.05	-188
2000	1775	159	6.08	43.27	49.35	0.12	-159
2001	1485	-131	48.45	0.90	49.35	0.98	131
2002	1650	34	25.85	23.50	49.35	0.52	-34
2003	1520	-96	47.95	1.40	49.35	0.97	96
2004	1483	-133	49.35	0.00	49.35	1.00	133
2005	1487	-129	49.35	0.00	49.35	1.00	129
2006	1648	32	26.45	22.90	49.35	0.54	-32
2007	1576	-40	37.35	12.00	49.35	0.76	40
2008	1649	33	25.90	23.45	49.35	0.52	-33
2009	1650	34	25.85	23.50	49.35	0.52	-34
2010	1570	-46	38.36	10.99	49.35	0.78	46
2011	1800	184	2.55	46.80	49.35	0.05	-184
MEAN	1587	-29	35.02	14.95		0.71	29



Photograph 50: Mt Genderme on 12 March 2011, resolution reduced to 200dpi.



Cirque glacier

GLACIER DATA		
Aspect	SW	
AREA	18.06	ha
Max Elev	1680	m
Min Elev	1310	m
Mean Elev	1495	m
Length	0.75	km
Elev Range	370	m
Gradient	0.49	

SNOWLINE DATA		
ELAo	1476	m
Max SL	1670	m, 1999
Min SL	1300	m, 1995
Mean SL	1460	m
SL Range	370	m
No. surveys	26	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE	DEPARTURE	AREAS		ACCUM.	MASS
	ELEVATION	FROM ELAo	ACCUM.	ABL.	TOTAL	AREA RATIO
	m	m	ha	ha	ha	(AAR)
ELAo	1476	0	9.26	8.80	18.06	0.51
1977						
1978	1480	4	8.96	9.10	18.06	0.50
1979						
1980						
1981	1408	-68	15.56	2.50	18.06	0.86
1982	1472	-4	9.35	8.71	18.06	0.52
1983	1344	-132	17.46	0.60	18.06	0.97
1984						
1985	1440	-36	12.77	5.29	18.06	0.71
1986						
1987	1454	-22	11.40	6.66	18.06	0.63
1988						
1989	1429	-47	13.96	4.10	18.06	0.77
1990						
1991						
1992						
1993	1360	-116	17.12	0.94	18.06	0.95
1994	1408	-68	15.60	2.46	18.06	0.86
1995	1300	-176	27.80	0.00	27.8	1.54
1996	1461	-15	10.54	7.52	18.06	0.58
1997	1321	-155	17.70	0.36	18.06	0.98
1998	1478	2	9.17	8.89	18.06	0.51
1999	1670	194	0.28	17.78	18.06	0.02
2000	1657	181	0.63	17.43	15.73	0.03
2001	1405	-71	15.72	2.34	18.06	0.87
2002	1613	137	2.07	12.52	14.59	0.11
2003	1376	-100	16.96	1.10	18.06	0.94
2004	1380	-96	17.06	1.00	18.06	0.94
2005	1400	-76	15.72	2.34	18.06	0.87
2006	1485	9	9.00	9.06	18.06	0.50
2007	1439	-37	13.95	5.10	18.06	0.77
2008	1580	104	3.26	14.80	18.06	0.18
2009	1600	124	2.46	15.60	18.06	0.14
2010	1408	-68	15.60	2.46	18.06	0.86
2011	1600	124	2.46	15.60	18.06	0.14
MEAN	1460	-16	11.64	6.70		0.64
						16



Photograph 51: Llawrenny Peaks on 12 March 2011, resolution reduced to 200dpi.



Mountain glacier

GLACIER DATA		
Aspect	S	
AREA	57.56	ha
Max Elev	1900	m
Min Elev	1420	m
Mean Elev	1660	m
Elev Range	480	m
Length	0.75	km
Gradient	0.64	

SNOWLINE DATA		
ELAo	1596	m
Max SL	1900	m, 1999
Min SL	1360	m, 1995
Mean SL	1584	m
SL Range	540	m
No. Surveys	27	

MEASUREMENTS Digitised values shaded

YEAR	SNOWLINE ELEVATION m	DEPARTURE FROM ELAo m	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1596	0	40.36	17.20	57.56	0.70	0
1977							
1978	1712	116	20.06	37.5	57.56	0.35	-116
1979							
1980	1545	-51	48.56	9	57.56	0.84	51
1981	1523	-73	51.23	6.33	57.56	0.89	73
1982	1565	-31	45.36	12.2	57.56	0.79	31
1983	1378	-218	63.9	0	63.9	1.11	218
1984							
1985	1524	-72	51.19	6.37	57.56	0.89	72
1986							
1987	1555	-41	46.86	10.7	57.56	0.81	41
1988							
1989	1525	-71	50.96	6.6	57.56	0.89	71
1990							
1991							
1992							
1993	1428	-168	60.7	0	60.7	1.05	168
1994	1478	-118	55.23	2.33	57.56	0.96	118
1995	1360	-236	65.03	0	65.03	1.13	236
1996	1470	-126	55.51	2.05	57.56	0.96	126
1997	1464	-132	56.75	0.81	57.56	0.99	132
1998	1682	86	25.14	32.42	57.56	0.44	-86
1999	1900	304	0	57.56	57.56	0.00	-304
2000	1803	207	3.28	54.28	57.56	0.06	-207
2001	1488	-108	54.56	3	57.56	0.95	108
2002	1790	194	5.36	52.2	57.56	0.09	-194
2003	1448	-148	56.56	1.00	57.56	0.98	148
2004	1465	-131	56.75	0.81	57.56	0.99	131
2005	1465	-131	56.75	0.81	57.56	0.99	131
2006	1700	104	20	37.56	57.56	0.35	-104
2007	1579	-17	43.36	14.2	57.56	0.75	17
2008	1715	119	19.56	38	57.56	0.34	-119
2009	1790	194	5.36	52.2	57.56	0.09	-194
2010	1555	-41	46.86	10.7	57.56	0.81	41
2011	1855	259	1.12	56.44	57.56	0.02	-259
MEAN	1584	-12	39.48	17.26	58.21	0.69	12



Photograph 52: Barrier Peak on 12 March 2011, resolution reduced to 200dpi.



GLACIER DATA			
	1977 to 1999	2002 on	
Aspect	E		
Permanent Ice Area	17.95	7.75	ha
Max Elev	1800	1770	m
Min Elev	1450	1510	m
Mean Elev	1625	1640	m
Elev Range	350	260	m
Length	0.5	0.43	km
Gradient	0.7	0.6	

SNOWLINE DATA		
ELAo	1563	m
Max SL	1770	m, 1978
Min SL	1400	m, 1995
Mean SL	1556	m
SL Range	370	m
No. Surveys	24	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	Permanent ice Outline Total ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo 1977 to 1999	1563	0	7.55	10.40	17.95	0.42	0
ELAo 1999 on	1600	0	3.25	4.50	7.75	0.42	0
1977							
1978	1700	137	0.83	17.12	17.95	0.05	-137
1979							
1980							
1981							
1982							
1983	1407	-156	17.95	0.00	17.95	1.00	156
1984							
1985	1526	-37	10.05	7.90	17.95	0.56	37
1986							
1987	1526	-37	10.05	7.90	17.95	0.56	37
1988							
1989	1537	-26	9.25	8.70	17.95	0.52	26
1990							
1991							
1992							
1993	1407	-156	31.25	0.00	17.95	1.00	156
1994	1512	-51	11.05	6.90	17.95	0.62	51
1995	1400	-163	32.65	0.00	17.95	1.00	163
1996	1498	-65	12.13	5.82	17.95	0.68	65
1997	1460	-103	17.95	0.00	17.95	1.00	103
1998	1612	49	4.73	13.22	17.95	0.26	-49
1999	1695	95	0.84	7.75	8.59	0.05	-95
2000	1700	100	0.83	7.27	8.1	0.10	-100
2001	1438	-162	17.60	0.36	8.1	1.00	162
2002	1770	170	0.34	7.41	7.75	0.04	-170
2003	1454	-146	15.85	0.00	7.75	1.00	146
2004	1470	-130	14.77	0.00	7.75	1.00	130
2005	1440	-160	17.55	0.00	7.75	1.00	160
2006	1670	70	1.80	5.95	7.75	0.23	-70
2007	1515	-85	11.05	0.00	7.75	1.00	85
2008	1650	50	2.60	5.15	7.75	0.34	-50
2009	1668	68	1.85	5.90	7.75	0.24	-68
2010	1510	-90	7.75	0.00	7.75	1.00	90
2011	1770	170	0.34	7.41	7.75	0.04	-170
MEAN	1556	-27	10.46	4.67		0.59	27



Photograph 53: Mt Irene on 12 March 2011, resolution reduced to 200dpi.

Patchy glacierette



GLACIER DATA		
Aspect	E	
AREA	12.78	ha
Max Elev	1700	m
Min Elev	1400	m
Mean Elev	1550	m
Elev Range	300	m
Length	0.5	km
Gradient	0.6	

SNOWLINE DATA		
ELAo	1515	m
Max SL	1690	m, 1999
Min SL	1350	m, 1995
Mean SL	1502	m
SL Range	340	m
No. Surveys	22	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1515	0	7.13	5.65	12.78	0.56	0
1977							
1978	1655	140	1.34	2.16	3.5	0.10	-140
1979					No visit		
1980					No visit		
1981					No visit		
1982					No visit		
1983					No visit		
1984					No visit		
1985					No visit		
1986					No visit		
1987					No visit		
1988					No visit		
1989	1545	30	5.83	6.95	12.78	0.46	-30
1990					No visit		
1991					No visit		
1992					No visit		
1993	1380	-135	16.20	0.00	16.2	1.27	135
1994	1420	-95	11.73	1.05	12.78	0.92	95
1995	1350	-165	19.34	0.00	19.34	1.51	165
1996	1425	-90	11.53	1.25	12.78	0.90	90
1997	1385	-130	15.65	0.00	15.65	1.22	130
1998	1445	-70	10.67	2.11	12.78	0.83	70
1999	1688	173	0.47	1.93	2.4	0.20	-173
2000					In cloud		
2001	1412	-103	12.38	0.4	12.78	0.97	103
2002	1690	175	0.42	1.76	2.18	0.19	-175
2003	1415	-100	11.88	0.9	12.78	0.93	100
2004	1405	-110	14.2	0	14.2	1.00	110
2005	1410	-105	12.38	0.4	12.7	0.97	105
2006	1665	150	1.3	1.7	3.0	0.43	-150
2007	1459	-56	8.3	2.7	11.0	0.65	56
2008	1670	155	1.1	1.4	2.5	0.45	-155
2009	1675	160	1.1	1.4	2.5	0.44	-160
2010	1440	-75	10.93	1.85	12.8	0.86	75
2011	SL above		0	2.5	2.5	0.00	
MEAN	1502	-13	8.34	1.47		0.75	13



Photograph 54: Merrie Range on 12 March 2011, resolution reduced to 200dpi.

Patchy glacierette




GLACIER DATA		
Aspect	SE	
AREA	9.99	ha
Max Elev	1600	m
Min Elev	1260	m
Mean Elev	1430	m
Elev Range	340	m
Length	0.5	km
Gradient	0.68	

SNOWLINE DATA		
ELAo	1380	m
Max SL	1575	m, 1999
Min SL	1220	m, 1993
Mean SL	1362	m
SL Range	355	m
No. Surveys	18	

MEASUREMENTS

Digitised values shaded

YEAR	SNOWLINE ELEVATION	DEPARTURE FROM ELAo	ACCUM. ha	AREAS ABL. ha	TOTAL ha	ACCUM. AREA RATIO (AAR)	MASS BALANCE INDEX
ELAo	1380	0	7.50	2.50	10	0.75	0
1977					No visit		
1978					No visit		
1979					No visit		
1980					No visit		
1981					No visit		
1982					No visit		
1983					No visit		
1984					No visit		
1985					No visit		
1986					No visit		
1987					No visit		
1988					No visit		
1989					No visit		
1990					No visit		
1991					No visit		
1992					No visit		
1993	1220	-160	15.79	0.00	15.79	1.00	160
1994	1381	1	8.95	1.04	7.60	1.18	-1
1995	1230	-150	14.90	0.00	14.90	1.00	150
1996	1302	-78	8.92	1.07	10.00	0.89	78
1997	1250	-130	13.30	0.00	13.30	1.00	130
1998	1382	2	7.43	2.56	9.99	0.74	-2
1999	1562	182	0.69	2.76	3.45	0.20	-182
2000					In cloud		
2001	1275	-105	9.40	3.60	13.00	0.72	105
2002	1575	195	0.46	2.27	2.73	0.17	-195
2003	1291	-89	9.10	3.00	12.10	0.75	89
2004	1270	-110	9.70	3.40	13.10	0.74	110
2005	1255	-125	9.75	3.55	13.30	0.73	125
2006	1410	30	6.60	0.60	7.20	0.92	-30
2007	1336	-44	8.30	1.90	10.20	0.81	44
2008	1430	50	5.80	1.00	6.80	0.85	-50
2009	1420	40	6.33	0.66	6.99	0.91	-40
2010	1380	0	7.38	0.34	7.72	0.96	0
2011	1550	170	0.90	2.60	3.50	0.26	-170
MEAN	1362	-18	8.61	1.78		0.78	18



Photograph 55: Caroline Peak on 12 March 2011, resolution reduced to 200dpi.